



CLINICAL PREVENTIVE GUIDELINE

Guideline Number: DHMP_PG1009

Effective Date: 11/1/2021

Guideline Subject: Pediatric, Adolescent and Adult
Immunization Criteria

Revision Date: 11/1/2022

Pages: 1 of 2

Gregg Kamas

10/1/2021

Quality Management Committee Chair

Date

I. PURPOSE:

Vaccines are among the most cost-effective clinical preventive services and are a core component of preventive services. Immunization and proper use of vaccines is an important defense against preventable infectious disease along with antibiotics, screening and testing guidelines and scientific improvements in the diagnosis of infectious disease-related health concerns (Healthy People 2020). The CDC recommends routine vaccination from birth through adulthood for lifetime protection against many diseases and infections.

This guideline will define the expected standards of health care for all eligible Denver Health Medical Plan (DHMP) pediatric, adolescent and adult members regarding immunizations within preventive health care services.

II. POPULATION:

All eligible members across all age groups for protection from infectious diseases through the use of vaccines. Use of vaccines reduces the risk of an individual transmitting infection to others, thereby conferring protection to the community.

III. GUIDELINE:

Vaccines will be recommended based on current nationally recognized vaccine-specific clinical care guidelines. These guidelines are based on the recommendations of the Advisory Committee on Immunization Practices (ACIP) of the Centers for Disease Control and Prevention (CDC) and the Colorado Department of Public Health and Environment (CDPHE). Further details may be accessed via https://www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fvaccines%2Fschedules%2Fhcp%2Fchild-adolescent.html CDC 2021 Updates/ Changes for Adolescents:

- The DTaP note was revised to include a “special situations” section containing information about the recommendation for use of DTaP in wound management.
- The Hib note was revised to indicate that for catch-up vaccination, no further doses are recommended if a previous dose was administered at age 15 months or older.
- The “birth dose” section of the HepB note contains additional text clarifying the recommendation for infants with birth weight of <2000 grams who have HBsAg-negative mothers.
- The HPV note was revised to include recommendations for interrupted schedules.
- The “special situations” section of the Influenza note has been revised for persons who have egg allergy with symptoms other than hives, and for situations where LAIV4 should not be used.
- The MenACWY note contains information about use of MenQuadfi, and the “special situations” section contains information about use of Menveo in infants who received dose 1 at age 3–6 months.
- The Tdap note was revised to include a “special situations” section containing information about the recommendation for use of Tdap in wound management.

NOTE:

This guideline is designed to assist providers by providing an analytical framework for the evaluation and treatment of patients, and is not intended either to replace a clinician's judgment or to establish a protocol for all patients with a particular condition.



CLINICAL PREVENTIVE GUIDELINE

Guideline Number: DHMP_PG1009

Effective Date: 11/1/2021

Guideline Subject: Pediatric, Adolescent and Adult
Immunization Criteria

Revision Date: 11/1/2022

Pages: 2 of 2

Quality Management Committee Chair

Date

- Although not included in spreadsheet Covid -19 Vaccine now recommended for those 18 and older for Moderna and Janssen, and 12 and older for Pfizer

Vaccination Schedules: as attachments below

1. Child and Adolescent Schedule (includes catch up schedule and special populations)
2. Adults, aged 19 years or older (includes special populations)

IV. ATTACHMENTS:

- A. Child and Adolescent Comprehensive Immunization Schedule (Includes catch-up schedule and special populations)
- B. Adult Immunization Schedule aged 19 years or older (Includes special populations)
- C. Parent Friendly Version Recommended Vaccinations for Children 0-6yrs
- D. Parent Friendly Version Recommended Vaccinations for Children 7-18 yrs
- E. Covid- 19 Vaccine Quick Reference Guide

V. REFERENCES:

- AAP Committee on Infectious Diseases. Recommended childhood and adolescent immunization schedules: United States, 2019. *Pediatrics* 2019 Feb 5; [e-pub].Retrived July 2021, from (<https://doi.org/10.1542/peds.2019-0065>)
- Centers for Disease Control and Prevention. (2021). Immunization Schedules. *National Center for Immunization and Respiratory Diseases*. Atlanta, GA, USA. Retrived July 2021, from <https://www.cdc.gov/vaccines/schedules/hcp/index.html>
- U.S. Department of Health and Human Services. (2021). *Healthy People 2020*. Retrived July 2021, From Immunization and Infectious Disease: <https://www.healthypeople.gov/2020/topics-objectives/topic/immunization-and-infectious-diseases>

NOTE:

This guideline is designed to assist providers by providing an analytical framework for the evaluation and treatment of patients, and is not intended either to replace a clinicians judgment or to establish a protocol for all patients with a particular condition.

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger

UNITED STATES
2021

Vaccines in the Child and Adolescent Immunization Schedule*

Vaccines	Abbreviations	Trade names
Diphtheria, tetanus, and acellular pertussis vaccine	DTaP	Daptacel® Infanrix®
Diphtheria, tetanus vaccine	DT	No trade name
<i>Haemophilus influenzae</i> type b vaccine	Hib (PRP-T) Hib (PRP-OMP)	ActHIB® Hiberix® PedvaxHIB®
Hepatitis A vaccine	HepA	Havrix® Vaqta®
Hepatitis B vaccine	HepB	Engerix-B® Recombivax HB®
Human papillomavirus vaccine	HPV	Gardasil 9®
Influenza vaccine (inactivated)	IIV	Multiple
Influenza vaccine (live, attenuated)	LAIV4	FluMist® Quadrivalent
Measles, mumps, and rubella vaccine	MMR	M-M-R II®
Meningococcal serogroups A, C, W, Y vaccine	MenACWY-D MenACWY-CRM MenACWY-TT	Menactra® Menveo® MenQuadfi®
Meningococcal serogroup B vaccine	MenB-4C MenB-FHbp	Bexsero® Trumenba®
Pneumococcal 13-valent conjugate vaccine	PCV13	Prevnar 13®
Pneumococcal 23-valent polysaccharide vaccine	PPSV23	Pneumovax 23®
Poliovirus vaccine (inactivated)	IPV	IPOL®
Rotavirus vaccine	RV1 RV5	Rotarix® RotaTeq®
Tetanus, diphtheria, and acellular pertussis vaccine	Tdap	Adacel® Boostrix®
Tetanus and diphtheria vaccine	Td	Tenivac® Tdvax™
Varicella vaccine	VAR	Varivax®

Combination vaccines (use combination vaccines instead of separate injections when appropriate)

DTaP, hepatitis B, and inactivated poliovirus vaccine	DTaP-HepB-IPV	Pediarix®
DTaP, inactivated poliovirus, and <i>Haemophilus influenzae</i> type b vaccine	DTaP-IPV/Hib	Pentacel®
DTaP and inactivated poliovirus vaccine	DTaP-IPV	Kinrix® Quadracel®
DTaP, inactivated poliovirus, <i>Haemophilus influenzae</i> type b, and hepatitis B vaccine	DTaP-IPV-Hib-HepB	Vaxelis®
Measles, mumps, rubella, and varicella vaccine	MMRV	ProQuad®

*Administer recommended vaccines if immunization history is incomplete or unknown. Do not restart or add doses to vaccine series for extended intervals between doses. When a vaccine is not administered at the recommended age, administer at a subsequent visit. The use of trade names is for identification purposes only and does not imply endorsement by the ACIP or CDC.

How to use the child/adolescent immunization schedule

- 1** Determine recommended vaccine by age (**Table 1**)
- 2** Determine recommended interval for catch-up vaccination (**Table 2**)
- 3** Assess need for additional recommended vaccines by medical condition and other indications (**Table 3**)
- 4** Review vaccine types, frequencies, intervals, and considerations for special situations (**Notes**)

Recommended by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/acip) and approved by the Centers for Disease Control and Prevention (www.cdc.gov), American Academy of Pediatrics (www.aap.org), American Academy of Family Physicians (www.aafp.org), American College of Obstetricians and Gynecologists (www.acog.org), American College of Nurse-Midwives (www.midwife.org), American Academy of Physician Assistants (www.aapa.org), and National Association of Pediatric Nurse Practitioners (www.napnap.org).

Report

- Suspected cases of reportable vaccine-preventable diseases or outbreaks to your state or local health department
- Clinically significant adverse events to the Vaccine Adverse Event Reporting System (VAERS) at www.vaers.hhs.gov or 800-822-7967



Download the CDC Vaccine Schedules App for providers at www.cdc.gov/vaccines/schedules/hcp/schedule-app.html.

Helpful information

- Complete ACIP recommendations: www.cdc.gov/vaccines/hcp/acip-recs/index.html
- *General Best Practice Guidelines for Immunization*: www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html
- Outbreak information (including case identification and outbreak response), see Manual for the Surveillance of Vaccine-Preventable Diseases: www.cdc.gov/vaccines/pubs/surv-manual
- ACIP Shared Clinical Decision-Making Recommendations www.cdc.gov/vaccines/acip/acip-scdm-faqs.html



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

Table 1

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2). School entry and adolescent vaccine age groups are shaded in gray.

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs	
Hepatitis B (HepB)	1 st dose	← 2 nd dose →		← 3 rd dose →														
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	See Notes													
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1 st dose	2 nd dose	3 rd dose	← 4 th dose →				5 th dose								
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	See Notes	← 3 rd or 4 th dose, See Notes →												
Pneumococcal conjugate (PCV13)			1 st dose	2 nd dose	3 rd dose	← 4 th dose →												
Inactivated poliovirus (IPV <18 yrs)			1 st dose	2 nd dose	← 3 rd dose →						4 th dose							
Influenza (IIV)											Annual vaccination 1 or 2 doses			Annual vaccination 1 dose only				
or											Annual vaccination 1 or 2 doses			Annual vaccination 1 dose only				
Influenza (LAIV4)											Annual vaccination 1 or 2 doses			Annual vaccination 1 dose only				
Measles, mumps, rubella (MMR)					See Notes	← 1 st dose →				2 nd dose								
Varicella (VAR)						← 1 st dose →				2 nd dose								
Hepatitis A (HepA)					See Notes	2-dose series, See Notes												
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)																		Tdap
Human papillomavirus (HPV)														*	See Notes			
Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos, MenACWY-TT ≥2years)			See Notes											1 st dose		2 nd dose		
Meningococcal B															See Notes			
Pneumococcal polysaccharide (PPSV23)												See Notes						

 Range of recommended ages for all children
 Range of recommended ages for catch-up immunization
 Range of recommended ages for certain high-risk groups
 Recommended based on shared clinical decision-making or *can be used in this age group
 No recommendation/not applicable

Table 2

Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 month Behind, United States, 2021

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with Table 1 and the notes that follow.**

Children age 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	4 weeks	8 weeks and at least 16 weeks after first dose. Minimum age for the final dose is 24 weeks.		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days.	4 weeks	4 weeks Maximum age for final dose is 8 months, 0 days.		
Diphtheria, tetanus, and acellular pertussis	6 weeks	4 weeks	4 weeks	6 months	6 months
<i>Haemophilus influenzae</i> type b	6 weeks	No further doses needed if first dose was administered at age 15 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose) if first dose was administered at age 12 through 14 months.	No further doses needed if previous dose was administered at age 15 months or older. 4 weeks if current age is younger than 12 months and first dose was administered at younger than age 7 months and at least 1 previous dose was PRP-T (ActHib, Pentacel, Hiberix) or unknown. 8 weeks and age 12 through 59 months (as final dose) if current age is younger than 12 months and first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months and first dose was administered before the 1 st birthday and second dose was administered at younger than 15 months; OR if both doses were PRP-OMP (PedvaxHIB, Comvax) and were administered before the 1 st birthday.	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before the 1 st birthday.	
Pneumococcal conjugate	6 weeks	No further doses needed for healthy children if first dose was administered at age 24 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose for healthy children) if first dose was administered at the 1 st birthday or after.	No further doses needed for healthy children if previous dose was administered at age 24 months or older. 4 weeks if current age is younger than 12 months and previous dose was administered at <7 months old. 8 weeks (as final dose for healthy children) if previous dose was administered between 7–11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was administered before age 12 months.	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before age 12 months or for children at high risk who received 3 doses at any age.	
Inactivated poliovirus	6 weeks	4 weeks	4 weeks if current age is <4 years. 6 months (as final dose) if current age is 4 years or older.	6 months (minimum age 4 years for final dose).	
Measles, mumps, rubella	12 months	4 weeks			
Varicella	12 months	3 months			
Hepatitis A	12 months	6 months			
Meningococcal ACWY	2 months MenACWY-CRM 9 months MenACWY-D 2 years MenACWY-TT	8 weeks	See Notes	See Notes	
Children and adolescents age 7 through 18 years					
Meningococcal ACWY	Not applicable (N/A)	8 weeks			
Tetanus, diphtheria; tetanus, diphtheria, and acellular pertussis	7 years	4 weeks	4 weeks if first dose of DTaP/DT was administered before the 1 st birthday. 6 months (as final dose) if first dose of DTaP/DT or Tdap/Td was administered at or after the 1 st birthday.	6 months if first dose of DTaP/DT was administered before the 1 st birthday.	
Human papillomavirus	9 years	Routine dosing intervals are recommended.			
Hepatitis A	N/A	6 months			
Hepatitis B	N/A	4 weeks	8 weeks and at least 16 weeks after first dose.		
Inactivated poliovirus	N/A	4 weeks	6 months A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years or if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	4 weeks			
Varicella	N/A	3 months if younger than age 13 years. 4 weeks if age 13 years or older.			

Table 3

Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2021

Always use this table in conjunction with Table 1 and the notes that follow.

VACCINE	INDICATION									
	Pregnancy	Immunocompromised status (excluding HIV infection)	HIV infection CD4+ count ¹		Kidney failure, end-stage renal disease, or on hemodialysis	Heart disease or chronic lung disease	CSF leak or cochlear implant	Asplenia or persistent complement deficiencies	Chronic liver disease	Diabetes
			<15% and total CD4 cell count of <200/mm ³	≥15% and total CD4 cell count of ≥200/mm ³						
Hepatitis B	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Rotavirus	Grey	Orange (SCID ²)	Orange	Orange	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Diphtheria, tetanus, and acellular pertussis (DTaP)	Grey	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
<i>Haemophilus influenzae</i> type b	Grey	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots
Pneumococcal conjugate	Grey	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots
Inactivated poliovirus	Orange	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Influenza (IIV)	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
or										
Influenza (LAIV4)	Red	Red	Red	Red	Orange (Asthma, wheezing: 2–4yrs ³)	Red	Red	Red	Orange	Orange
Measles, mumps, rubella	Red (*)	Red	Red	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Varicella	Red (*)	Red	Red	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Hepatitis A	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Tetanus, diphtheria, and acellular pertussis (Tdap)	Yellow with dots	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Human papillomavirus	Red (*)	Yellow with dots	Yellow with dots	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Meningococcal ACWY	Yellow	Yellow	Yellow with dots	Yellow	Yellow	Yellow	Yellow	Yellow with dots	Yellow	Yellow
Meningococcal B	Orange	Purple	Purple	Purple	Purple	Purple	Purple	Yellow with dots	Purple	Purple
Pneumococcal polysaccharide	Purple	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots	Yellow with dots

Yellow Vaccination according to the routine schedule recommended
 Purple Recommended for persons with an additional risk factor for which the vaccine would be indicated
 Yellow with dots Vaccination is recommended, and additional doses may be necessary based on medical condition. See Notes.
 Red Not recommended/contraindicated—vaccine should not be administered.
 Orange Precaution—vaccine might be indicated if benefit of protection outweighs risk of adverse reaction
 Grey No recommendation/not applicable

1 For additional information regarding HIV laboratory parameters and use of live vaccines, see the *General Best Practice Guidelines for Immunization*, "Altered Immunocompetence," at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html and Table 4-1 (footnote D) at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html.

2 Severe Combined Immunodeficiency

3 LAIV4 contraindicated for children 2–4 years of age with asthma or wheezing during the preceding 12 months

For vaccination recommendations for persons ages 19 years or older, see the Recommended Adult Immunization Schedule, 2021.

Additional information

COVID-19 Vaccination

ACIP recommends use of COVID-19 vaccines within the scope of the Emergency Use Authorization or Biologics License Application for the particular vaccine. Interim ACIP recommendations for the use of COVID-19 vaccines can be found at www.cdc.gov/vaccines/hcp/acip-recs/.

- Consult relevant ACIP statements for detailed recommendations at www.cdc.gov/vaccines/hcp/acip-recs/index.html.
- For information on contraindications and precautions for the use of a vaccine, consult the *General Best Practice Guidelines for Immunization* at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html and relevant ACIP statements at www.cdc.gov/vaccines/hcp/acip-recs/index.html.
- For calculating intervals between doses, 4 weeks = 28 days. Intervals of ≥ 4 months are determined by calendar months.
- Within a number range (e.g., 12–18), a dash (–) should be read as “through.”
- Vaccine doses administered ≤ 4 days before the minimum age or interval are considered valid. Doses of any vaccine administered ≥ 5 days earlier than the minimum age or minimum interval should not be counted as valid and should be repeated as age appropriate. **The repeat dose should be spaced after the invalid dose by the recommended minimum interval.** For further details, see Table 3-1, Recommended and minimum ages and intervals between vaccine doses, in *General Best Practice Guidelines for Immunization* at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/timing.html.
- Information on travel vaccination requirements and recommendations is available at www.cdc.gov/travel/.
- For vaccination of persons with immunodeficiencies, see Table 8-1, Vaccination of persons with primary and secondary immunodeficiencies, in *General Best Practice Guidelines for Immunization* at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html, and Immunization in Special Clinical Circumstances (In: Kimberlin DW, Brady MT, Jackson MA, Long SS, eds. *Red Book: 2018 Report of the Committee on Infectious Diseases*. 31st ed. Itasca, IL: American Academy of Pediatrics; 2018:67–111).
- For information about vaccination in the setting of a vaccine-preventable disease outbreak, contact your state or local health department.
- The National Vaccine Injury Compensation Program (VICP) is a no-fault alternative to the traditional legal system for resolving vaccine injury claims. All routine child and adolescent vaccines are covered by VICP except for pneumococcal polysaccharide vaccine (PPSV23). For more information, see www.hrsa.gov/vaccinecompensation/index.html.

Diphtheria, tetanus, and pertussis (DTaP) vaccination (minimum age: 6 weeks [4 years for Kinrix or Quadacel])

Routine vaccination

- 5-dose series at 2, 4, 6, 15–18 months, 4–6 years
 - **Prospectively:** Dose 4 may be administered as early as age 12 months if at least 6 months have elapsed since dose 3.
 - **Retrospectively:** A 4th dose that was inadvertently administered as early as age 12 months may be counted if at least 4 months have elapsed since dose 3.

Catch-up vaccination

- Dose 5 is not necessary if dose 4 was administered at age 4 years or older and at least 6 months after dose 3.
- For other catch-up guidance, see Table 2.

Special situations

- Wound management in children less than age 7 years with history of 3 or more doses of tetanus-toxoid-containing vaccine: For all wounds except clean and minor wounds, administer DTaP if more than 5 years since last dose of tetanus-toxoid-containing vaccine. For detailed information, see www.cdc.gov/mmwr/volumes/67/rr/rr6702a1.htm.

Haemophilus influenzae type b vaccination (minimum age: 6 weeks)

Routine vaccination

- **ActHIB, Hiberix, or Pentacel:** 4-dose series at 2, 4, 6, 12–15 months
- **PedvaxHIB:** 3-dose series at 2, 4, 12–15 months

Catch-up vaccination

- **Dose 1 at age 7–11 months:** Administer dose 2 at least 4 weeks later and dose 3 (final dose) at age 12–15 months or 8 weeks after dose 2 (whichever is later).
- **Dose 1 at age 12–14 months:** Administer dose 2 (final dose) at least 8 weeks after dose 1.
- **Dose 1 before age 12 months and dose 2 before age 15 months:** Administer dose 3 (final dose) 8 weeks after dose 2.
- **2 doses of PedvaxHIB before age 12 months:** Administer dose 3 (final dose) at 12–59 months and at least 8 weeks after dose 2.
- **1 dose administered at age 15 months or older:** No further doses needed
- **Unvaccinated at age 15–59 months:** Administer 1 dose.
- **Previously unvaccinated children age 60 months or older who are not considered high risk:** Do not require catch-up vaccination
- For other catch-up guidance, see Table 2.

Special situations

• Chemotherapy or radiation treatment:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Doses administered within 14 days of starting therapy or during therapy should be repeated at least 3 months after therapy completion.

• Hematopoietic stem cell transplant (HSCT):

- 3-dose series 4 weeks apart starting 6 to 12 months after successful transplant, regardless of Hib vaccination history

• Anatomic or functional asplenia (including sickle cell disease):

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Unvaccinated* persons age 5 years or older

- 1 dose

• Elective splenectomy:

Unvaccinated* persons age 15 months or older

- 1 dose (preferably at least 14 days before procedure)

• HIV infection:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

Unvaccinated* persons age 5–18 years

- 1 dose

• Immunoglobulin deficiency, early component complement deficiency:

12–59 months

- Unvaccinated or only 1 dose before age 12 months: 2 doses, 8 weeks apart
- 2 or more doses before age 12 months: 1 dose at least 8 weeks after previous dose

*Unvaccinated = Less than routine series (through age 14 months) OR no doses (age 15 months or older)

Hepatitis A vaccination

(minimum age: 12 months for routine vaccination)

Routine vaccination

- 2-dose series (minimum interval: 6 months) beginning at age 12 months

Catch-up vaccination

- Unvaccinated persons through age 18 years should complete a 2-dose series (minimum interval: 6 months).
- Persons who previously received 1 dose at age 12 months or older should receive dose 2 at least 6 months after dose 1.
- Adolescents age 18 years or older may receive the combined HepA and HepB vaccine, **Twinrix**[®], as a 3-dose series (0, 1, and 6 months) or 4-dose series (3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months).

International travel

- Persons traveling to or working in countries with high or intermediate endemic hepatitis A (www.cdc.gov/travel/):
 - **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2 doses, separated by at least 6 months, between age 12–23 months.
 - **Unvaccinated age 12 months or older:** Administer dose 1 as soon as travel is considered.

Hepatitis B vaccination

(minimum age: birth)

Birth dose (monovalent HepB vaccine only)

- **Mother is HBsAg-negative:** 1 dose within 24 hours of birth for all medically stable infants $\geq 2,000$ grams. Infants $< 2,000$ grams: Administer 1 dose at chronological age 1 month or hospital discharge (whichever is earlier and even if weight is still $< 2,000$ grams).
- **Mother is HBsAg-positive:**
 - Administer **HepB vaccine** and **hepatitis B immune globulin (HBIG)** (in separate limbs) within 12 hours of birth, regardless of birth weight. For infants $< 2,000$ grams, administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
 - Test for HBsAg and anti-HBs at age 9–12 months. If HepB series is delayed, test 1–2 months after final dose.
- **Mother's HBsAg status is unknown:**
 - Administer **HepB vaccine** within 12 hours of birth, regardless of birth weight.
 - For infants $< 2,000$ grams, administer **HBIG** in addition to HepB vaccine (in separate limbs) within 12 hours of birth. Administer 3 additional doses of vaccine (total of 4 doses) beginning at age 1 month.
 - Determine mother's HBsAg status as soon as possible. If mother is HBsAg-positive, administer **HBIG** to infants $\geq 2,000$ grams as soon as possible, but no later than 7 days of age.

Routine series

- 3-dose series at 0, 1–2, 6–18 months (use monovalent HepB vaccine for doses administered before age 6 weeks)
- Infants who did not receive a birth dose should begin the series as soon as feasible (see Table 2).
- Administration of **4 doses** is permitted when a combination vaccine containing HepB is used after the birth dose.

- **Minimum age** for the final (3rd or 4th) dose: 24 weeks
- **Minimum intervals:** dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 8 weeks / dose 1 to dose 3: 16 weeks (when 4 doses are administered, substitute "dose 4" for "dose 3" in these calculations)

Catch-up vaccination

- Unvaccinated persons should complete a 3-dose series at 0, 1–2, 6 months.
- Adolescents age 11–15 years may use an alternative 2-dose schedule with at least 4 months between doses (adult formulation **Recombivax HB** only).
- Adolescents age 18 years or older may receive a 2-dose series of HepB (**Heplisav-B**[®]) at least 4 weeks apart.
- Adolescents age 18 years or older may receive the combined HepA and HepB vaccine, **Twinrix**, as a 3-dose series (0, 1, and 6 months) or 4-dose series (3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months).
- For other catch-up guidance, see Table 2.

Special situations

- Revaccination is not generally recommended for persons with a normal immune status who were vaccinated as infants, children, adolescents, or adults.
- **Revaccination** may be recommended for certain populations, including:
 - **Infants born to HBsAg-positive mothers**
 - **Hemodialysis patients**
 - **Other immunocompromised persons**
- For detailed revaccination recommendations, see www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/hepb.html.

Human papillomavirus vaccination

(minimum age: 9 years)

Routine and catch-up vaccination

- HPV vaccination routinely recommended at **age 11–12 years (can start at age 9 years)** and catch-up HPV vaccination recommended for all persons through age 18 years if not adequately vaccinated
- 2- or 3-dose series depending on age at initial vaccination:
 - **Age 9–14 years at initial vaccination:** 2-dose series at 0, 6–12 months (minimum interval: 5 months; repeat dose if administered too soon)
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2 months, 6 months (minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months; repeat dose if administered too soon)
- **Interrupted schedules:** If vaccination schedule is interrupted, the series does not need to be restarted.
- No additional dose recommended after completing series with recommended dosing intervals using any HPV vaccine.

Special situations

- **Immunocompromising conditions, including HIV infection:** 3-dose series as above
- **History of sexual abuse or assault:** Start at age 9 years.
- **Pregnancy:** HPV vaccination not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination

Influenza vaccination

(minimum age: 6 months [IIV], 2 years [LAIV4], 18 years [recombinant influenza vaccine, RIV4])

Routine vaccination

- Use any influenza vaccine appropriate for age and health status annually:
 - 2 doses, separated by at least 4 weeks, for **children age 6 months–8 years** who have received fewer than 2 influenza vaccine doses before July 1, 2020, or whose influenza vaccination history is unknown (administer dose 2 even if the child turns 9 between receipt of dose 1 and dose 2)
 - 1 dose for **children age 6 months–8 years** who have received at least 2 influenza vaccine doses before July 1, 2020
 - 1 dose for **all persons age 9 years or older**
- For the 2021–22 season, see the 2021–22 ACIP influenza vaccine recommendations.

Special situations

- **Egg allergy, hives only:** Any influenza vaccine appropriate for age and health status annually
- **Egg allergy with symptoms other than hives** (e.g., angioedema, respiratory distress, need for emergency medical services or epinephrine): Any influenza vaccine appropriate for age and health status annually. If using an influenza vaccine other than Flublok or Flucelvax, administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions.
- Severe allergic reactions to vaccines can occur even in the absence of a history of previous allergic reaction. All vaccination providers should be familiar with the office emergency plan and certified in cardiopulmonary resuscitation.
- A previous severe allergic reaction to influenza vaccine is a contraindication to future receipt of any influenza vaccine.
- **LAIV4 should not be used** in persons with the following conditions or situations:
 - History of severe allergic reaction to a previous dose of any influenza vaccine or to any vaccine component (excluding egg, see details above)
 - Receiving aspirin or salicylate-containing medications
 - Age 2–4 years with history of asthma or wheezing
 - Immunocompromised due to any cause (including medications and HIV infection)
 - Anatomic or functional asplenia
 - Close contacts or caregivers of severely immunosuppressed persons who require a protected environment
 - Pregnancy
 - Cochlear implant
 - Cerebrospinal fluid-oro-pharyngeal communication
 - Children less than age 2 years
 - Received influenza antiviral medications oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days

Notes

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021

Measles, mumps, and rubella vaccination (minimum age: 12 months for routine vaccination)

Routine vaccination

- 2-dose series at 12–15 months, 4–6 years
- Dose 2 may be administered as early as 4 weeks after dose 1.

Catch-up vaccination

- Unvaccinated children and adolescents: 2-dose series at least 4 weeks apart
- The maximum age for use of MMRV is 12 years.

Special situations

International travel

- **Infants age 6–11 months:** 1 dose before departure; revaccinate with 2-dose series at age 12–15 months (12 months for children in high-risk areas) and dose 2 as early as 4 weeks later.
- **Unvaccinated children age 12 months or older:** 2-dose series at least 4 weeks apart before departure

Meningococcal serogroup A,C,W,Y vaccination (minimum age: 2 months [MenACWY-CRM, Menveo], 9 months [MenACWY-D, Menactra], 2 years [MenACWY-TT, MenQuadfi])

Routine vaccination

- 2-dose series at 11–12 years, 16 years

Catch-up vaccination

- Age 13–15 years: 1 dose now and booster at age 16–18 years (minimum interval: 8 weeks)
- Age 16–18 years: 1 dose

Special situations

Anatomic or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- **Menveo**
 - Dose 1 at age 8 weeks: 4-dose series at 2, 4, 6, 12 months
 - Dose 1 at age 3–6 months: 3- or 4- dose series (dose 2 [and dose 3 if applicable] at least 8 weeks after previous dose until a dose is received at age 7 months or older, followed by an additional dose at least 12 weeks later and after age 12 months)
 - Dose 1 at age 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - Dose 1 at age 24 months or older: 2-dose series at least 8 weeks apart
- **Menactra**
 - **Persistent complement component deficiency or complement inhibitor use:**
 - Age 9–23 months: 2-dose series at least 12 weeks apart
 - Age 24 months or older: 2-dose series at least 8 weeks apart
 - **Anatomic or functional asplenia, sickle cell disease, or HIV infection:**
 - Age 9–23 months: Not recommended
 - Age 24 months or older: 2-dose series at least 8 weeks apart
 - **Menactra** must be administered at least 4 weeks after completion of PCV13 series.

- **MenQuadfi**

- Dose 1 at age 24 months or older: 2-dose series at least 8 weeks apart

Travel in countries with hyperendemic or epidemic meningococcal disease, including countries in the African meningitis belt or during the Hajj (www.cdc.gov/travel/):

- Children less than age 24 months:
 - **Menveo (age 2–23 months)**
 - Dose 1 at age 8 weeks: 4-dose series at 2, 4, 6, 12 months
 - Dose 1 at age 3–6 months: 3- or 4- dose series (dose 2 [and dose 3 if applicable] at least 8 weeks after previous dose until a dose is received at age 7 months or older, followed by an additional dose at least 12 weeks later and after age 12 months)
 - Dose 1 at age 7–23 months: 2-dose series (dose 2 at least 12 weeks after dose 1 and after age 12 months)
 - **Menactra (age 9–23 months)**
 - 2-dose series (dose 2 at least 12 weeks after dose 1; dose 2 may be administered as early as 8 weeks after dose 1 in travelers)
- Children age 2 years or older: 1 dose Menveo, Menactra, or MenQuadfi

First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) or military recruits:

- 1 dose **Menveo, Menactra, or MenQuadfi**
- Adolescent vaccination of children who received MenACWY prior to age 10 years:**
- **Children for whom boosters are recommended** because of an ongoing increased risk of meningococcal disease (e.g., those with complement deficiency, HIV, or asplenia): Follow the booster schedule for persons at increased risk.
 - **Children for whom boosters are not recommended** (e.g., a healthy child who received a single dose for travel to a country where meningococcal disease is endemic): Administer MenACWY according to the recommended adolescent schedule with dose 1 at age 11–12 years and dose 2 at age 16 years.

Note: **Menactra** should be administered either before or at the same time as DTaP. For MenACWY **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm.

Meningococcal serogroup B vaccination (minimum age: 10 years [MenB-4C, Bexsero; MenB-FHbp, Trumenba])

Shared clinical decision-making

- **Adolescents not at increased risk** age 16–23 years (preferred age 16–18 years) based on shared clinical decision-making:
 - **Bexsero:** 2-dose series at least 1 month apart
 - **Trumenba:** 2-dose series at least 6 months apart; if dose 2 is administered earlier than 6 months, administer a 3rd dose at least 4 months after dose 2.

Special situations

Anatomic or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:

- **Bexsero:** 2-dose series at least 1 month apart
 - **Trumenba:** 3-dose series at 0, 1–2, 6 months
- Bexsero** and **Trumenba** are not interchangeable; the same product should be used for all doses in a series. For MenB **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm.

Pneumococcal vaccination

(minimum age: 6 weeks [PCV13], 2 years [PPSV23])

Routine vaccination with PCV13

- 4-dose series at 2, 4, 6, 12–15 months

Catch-up vaccination with PCV13

- 1 dose for healthy children age 24–59 months with any incomplete* PCV13 series
- For other catch-up guidance, see Table 2.

Special situations

Underlying conditions below: When both PCV13 and PPSV23 are indicated, administer PCV13 first. PCV13 and PPSV23 should not be administered during same visit.

Chronic heart disease (particularly cyanotic congenital heart disease and cardiac failure); chronic lung disease (including asthma treated with high-dose, oral corticosteroids); diabetes mellitus:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after completing all recommended PCV13 doses)

Age 6–18 years

- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after completing all recommended PCV13 doses)

Cerebrospinal fluid leak, cochlear implant:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

Age 6–18 years

- No history of either PCV13 or PPSV23: 1 dose PCV13, 1 dose PPSV23 at least 8 weeks later
- Any PCV13 but no PPSV23: 1 dose PPSV23 at least 8 weeks after the most recent dose of PCV13
- PPSV23 but no PCV13: 1 dose PCV13 at least 8 weeks after the most recent dose of PPSV23

Sickle cell disease and other hemoglobinopathies; anatomic or functional asplenia; congenital or acquired immunodeficiency; HIV infection; chronic renal failure; nephrotic syndrome; malignant neoplasms, leukemias, lymphomas, Hodgkin disease, and other diseases associated with treatment with immunosuppressive drugs or radiation therapy; solid organ transplantation; multiple myeloma:

Age 2–5 years

- Any incomplete* series with:
 - 3 PCV13 doses: 1 dose PCV13 (at least 8 weeks after any prior PCV13 dose)
 - Less than 3 PCV13 doses: 2 doses PCV13 (8 weeks after the most recent dose and administered 8 weeks apart)
- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose) and a 2nd dose of PPSV23 5 years later

Age 6–18 years

- No history of either PCV13 or PPSV23: 1 dose PCV13, 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after PCV13 and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- Any PCV13 but no PPSV23: 2 doses PPSV23 (dose 1 of PPSV23 administered 8 weeks after the most recent dose of PCV13 and dose 2 of PPSV23 administered at least 5 years after dose 1 of PPSV23)
- PPSV23 but no PCV13: 1 dose PCV13 at least 8 weeks after the most recent PPSV23 dose and a 2nd dose of PPSV23 administered 5 years after dose 1 of PPSV23 and at least 8 weeks after a dose of PCV13

Chronic liver disease, alcoholism:

Age 6–18 years

- No history of PPSV23: 1 dose PPSV23 (at least 8 weeks after any prior PCV13 dose)

**Incomplete series* = Not having received all doses in either the recommended series or an age-appropriate catch-up series. See Tables 8, 9, and 11 in the ACIP pneumococcal vaccine recommendations (www.cdc.gov/mmwr/pdf/rr/rr5911.pdf) for complete schedule details.

Poliovirus vaccination (minimum age: 6 weeks)

Routine vaccination

- 4-dose series at ages 2, 4, 6–18 months, 4–6 years; administer the final dose on or after age 4 years and at least 6 months after the previous dose.
- 4 or more doses of IPV can be administered before age 4 years when a combination vaccine containing IPV is used. However, a dose is still recommended on or after age 4 years and at least 6 months after the previous dose.

Catch-up vaccination

- In the first 6 months of life, use minimum ages and intervals only for travel to a polio-endemic region or during an outbreak.
- IPV is not routinely recommended for U.S. residents age 18 years or older.

Series containing oral polio vaccine (OPV), either mixed OPV-IPV or OPV-only series:

- Total number of doses needed to complete the series is the same as that recommended for the U.S. IPV schedule. See www.cdc.gov/mmwr/volumes/66/wr/mm6601a6.htm?s_cid=mm6601a6_w.
- Only trivalent OPV (tOPV) counts toward the U.S. vaccination requirements.
 - Doses of OPV administered before April 1, 2016, should be counted (unless specifically noted as administered during a campaign).
 - Doses of OPV administered on or after April 1, 2016, should not be counted.
 - For guidance to assess doses documented as “OPV,” see www.cdc.gov/mmwr/volumes/66/wr/mm6606a7.htm?s_cid=mm6606a7_w.
- For other catch-up guidance, see Table 2.

Rotavirus vaccination (minimum age: 6 weeks)

Routine vaccination

- **Rotarix:** 2-dose series at 2 and 4 months
- **RotaTeq:** 3-dose series at 2, 4, and 6 months
- If any dose in the series is either **RotaTeq** or unknown, default to 3-dose series.

Catch-up vaccination

- Do not start the series on or after age 15 weeks, 0 days.
- The maximum age for the final dose is 8 months, 0 days.
- For other catch-up guidance, see Table 2.

Tetanus, diphtheria, and pertussis (Tdap) vaccination

(minimum age: 11 years for routine vaccination, 7 years for catch-up vaccination)

Routine vaccination

- **Adolescents age 11–12 years:** 1 dose Tdap
- **Pregnancy:** 1 dose Tdap during each pregnancy, preferably in early part of gestational weeks 27–36
- Tdap may be administered regardless of the interval since the last tetanus- and diphtheria-toxoid-containing vaccine.

Catch-up vaccination

- **Adolescents age 13–18 years who have not received Tdap:** 1 dose Tdap, then Td or Tdap booster every 10 years
- **Persons age 7–18 years not fully vaccinated* with DTaP:** 1 dose Tdap as part of the catch-up series (preferably the first dose); if additional doses are needed, use Td or Tdap.
- **Tdap administered at age 7–10 years:**
 - **Children age 7–9 years** who receive Tdap should receive the routine Tdap dose at age 11–12 years.
 - **Children age 10 years** who receive Tdap do not need the routine Tdap dose at age 11–12 years.
- **DTaP inadvertently administered on or after age 7 years:**
 - **Children age 7–9 years:** DTaP may count as part of catch-up series. Administer routine Tdap dose at age 11–12 years.
 - **Children age 10–18 years:** Count dose of DTaP as the adolescent Tdap booster.
- For other catch-up guidance, see Table 2.

Special situations

- **Wound management** in persons age 7 years or older with history of 3 or more doses of tetanus-toxoid-containing vaccine: For clean and minor wounds, administer Tdap or Td if more than 10 years since last dose of tetanus-toxoid-containing vaccine; for all other wounds, administer Tdap or Td if more than 5 years since last dose of tetanus-toxoid-containing vaccine. Tdap is preferred for persons age 11 years or older who have not previously received Tdap or whose Tdap history is unknown. If a tetanus-toxoid-containing vaccine is indicated for a pregnant adolescent, use Tdap.
- For detailed information, see www.cdc.gov/mmwr/volumes/69/wr/mm6903a5.htm.

**Fully vaccinated* = 5 valid doses of DTaP OR 4 valid doses of DTaP if dose 4 was administered at age 4 years or older

Varicella vaccination (minimum age: 12 months)

Routine vaccination

- 2-dose series at 12–15 months, 4–6 years
- Dose 2 may be administered as early as 3 months after dose 1 (a dose administered after a 4-week interval may be counted).

Catch-up vaccination

- Ensure persons age 7–18 years without evidence of immunity (see *MMWR* at www.cdc.gov/mmwr/pdf/rr/rr5604.pdf) have a 2-dose series:
 - **Age 7–12 years:** routine interval: 3 months (a dose administered after a 4-week interval may be counted)
 - **Age 13 years and older:** routine interval: 4–8 weeks (minimum interval: 4 weeks)
 - The maximum age for use of MMRV is 12 years.

Recommended Adult Immunization Schedule for ages 19 years or older

UNITED STATES
2021

How to use the adult immunization schedule

- 1** Determine recommended vaccinations by age (**Table 1**)
- 2** Assess need for additional recommended vaccinations by medical condition and other indications (**Table 2**)
- 3** Review vaccine types, frequencies, and intervals and considerations for special situations (**Notes**)

Recommended by the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/acip) and approved by the Centers for Disease Control and Prevention (www.cdc.gov), American College of Physicians (www.acponline.org), American Academy of Family Physicians (www.aafp.org), American College of Obstetricians and Gynecologists (www.acog.org), American College of Nurse-Midwives (www.midwife.org), and American Academy of Physician Assistants (www.aapa.org).

Vaccines in the Adult Immunization Schedule*

Vaccines	Abbreviations	Trade names
<i>Haemophilus influenzae</i> type b vaccine	Hib	ActHIB® Hiberix® PedvaxHIB®
Hepatitis A vaccine	HepA	Havrix® Vaqta®
Hepatitis A and hepatitis B vaccine	HepA-HepB	Twinrix®
Hepatitis B vaccine	HepB	Engerix-B® Recombivax HB® Hepelisav-B®
Human papillomavirus vaccine	HPV	Gardasil 9®
Influenza vaccine (inactivated)	IIV	Many brands
Influenza vaccine (live, attenuated)	LAIV4	FluMist® Quadrivalent
Influenza vaccine (recombinant)	RIV4	Flublok® Quadrivalent
Measles, mumps, and rubella vaccine	MMR	M-M-R II®
Meningococcal serogroups A, C, W, Y vaccine	MenACWY-D MenACWY-CRM MenACWY-TT	Menactra® Menveo® MenQuadfi®
Meningococcal serogroup B vaccine	MenB-4C MenB-FHbp	Bexsero® Trumenba®
Pneumococcal 13-valent conjugate vaccine	PCV13	Prevnar 13®
Pneumococcal 23-valent polysaccharide vaccine	PPSV23	Pneumovax 23®
Tetanus and diphtheria toxoids	Td	Tenivac® Tdvax™
Tetanus and diphtheria toxoids and acellular pertussis vaccine	Tdap	Adacel® Boostrix®
Varicella vaccine	VAR	Varivax®
Zoster vaccine, recombinant	RZV	Shingrix

*Administer recommended vaccines if vaccination history is incomplete or unknown. Do not restart or add doses to vaccine series if there are extended intervals between doses. The use of trade names is for identification purposes only and does not imply endorsement by the ACIP or CDC.

Report

- Suspected cases of reportable vaccine-preventable diseases or outbreaks to the local or state health department
- Clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System at www.vaers.hhs.gov or 800-822-7967

Injury claims

All vaccines included in the adult immunization schedule except pneumococcal 23-valent polysaccharide (PPSV23) and zoster (RZV) vaccines are covered by the Vaccine Injury Compensation Program. Information on how to file a vaccine injury claim is available at www.hrsa.gov/vaccinecompensation.

Questions or comments

Contact www.cdc.gov/cdc-info or 800-CDC-INFO (800-232-4636), in English or Spanish, 8 a.m.–8 p.m. ET, Monday through Friday, excluding holidays.



Download the CDC Vaccine Schedules app for providers at www.cdc.gov/vaccines/schedules/hcp/schedule-app.html.

Helpful information

- Complete ACIP recommendations: www.cdc.gov/vaccines/hcp/acip-recs/index.html
- *General Best Practice Guidelines for Immunization* (including contraindications and precautions): www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html
- Vaccine information statements: www.cdc.gov/vaccines/hcp/vis/index.html
- Manual for the Surveillance of Vaccine-Preventable Diseases (including case identification and outbreak response): www.cdc.gov/vaccines/pubs/surv-manual
- Travel vaccine recommendations: www.cdc.gov/travel
- Recommended Child and Adolescent Immunization Schedule, United States, 2021: www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html
- ACIP Shared Clinical Decision-Making Recommendations www.cdc.gov/vaccines/acip/acip-scdm-faqs.html



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2021

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
Influenza inactivated (IIV) or Influenza recombinant (RIV4) ^{or}	1 dose annually			
Influenza live, attenuated (LAIV4)				
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			
Varicella (VAR)	2 doses (if born in 1980 or later)		2 doses	
Zoster recombinant (RZV)			2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal conjugate (PCV13)	1 dose			1 dose
Pneumococcal polysaccharide (PPSV23)	1 or 2 doses depending on indication			1 dose
Hepatitis A (HepA)	2 or 3 doses depending on vaccine			
Hepatitis B (HepB)	2 or 3 doses depending on vaccine			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations			
	19 through 23 years			
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection

Recommended vaccination for adults with an additional risk factor or another indication

Recommended vaccination based on shared clinical decision-making

No recommendation/ Not applicable

Table 2 Recommended Adult Immunization Schedule by Medical Condition and Other Indications, United States, 2021

Vaccine	Pregnancy	Immuno-compromised (excluding HIV infection)	HIV infection CD4 count		Asplenia, complement deficiencies	End-stage renal disease; or on hemodialysis	Heart or lung disease, alcoholism ¹	Chronic liver disease	Diabetes	Health care personnel ²	Men who have sex with men
			<200 mm ³	≥200 mm ³							
IIV or RIV4 or	1 dose annually										
LAIV4	Not Recommended					Precaution			1 dose annually or		
Tdap or Td	1 dose Tdap each pregnancy	1 dose Tdap, then Td or Tdap booster every 10 years									
MMR	Not Recommended*	Not Recommended	1 or 2 doses depending on indication								
VAR	Not Recommended*	Not Recommended		2 doses							
RZV			2 doses at age ≥50 years								
HPV	Not Recommended*	3 doses through age 26 years		2 or 3 doses through age 26 years depending on age at initial vaccination or condition							
PCV13		1 dose									
PPSV23		1, 2, or 3 doses depending on age and indication									
HepA				2 or 3 doses depending on vaccine							
HepB				2, 3, or 4 doses depending on vaccine or condition				<60 years			
								≥60 years			
MenACWY	1 or 2 doses depending on indication, see notes for booster recommendations										
MenB	Precaution	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations									
Hib		3 doses HSCT ³ recipients only		1 dose							

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of past infection

 Recommended vaccination for adults with an additional risk factor or another indication

 Precaution—vaccination might be indicated if benefit of protection outweighs risk of adverse reaction

 Recommended vaccination based on shared clinical decision-making

 Not recommended/contraindicated—vaccine should not be administered.

 No recommendation/Not applicable

*Vaccinate after pregnancy.

1. Precaution for LAIV4 does not apply to alcoholism. 2. See notes for influenza; hepatitis B; measles, mumps, and rubella; and varicella vaccinations. 3. Hematopoietic stem cell transplant.

For vaccine recommendations for persons 18 years of age or younger, see the Recommended Child/Adolescent Immunization Schedule.

Additional Information

COVID-19 Vaccination

ACIP recommends use of COVID-19 vaccines within the scope of the Emergency Use Authorization or Biologics License Application for the particular vaccine. Interim ACIP recommendations for the use of COVID-19 vaccines can be found at www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/covid-19.html

Haemophilus influenzae type b vaccination

Special situations

- **Anatomical or functional asplenia (including sickle cell disease):** 1 dose if previously did not receive Hib; if elective splenectomy, 1 dose, preferably at least 14 days before splenectomy
- **Hematopoietic stem cell transplant (HSCT):** 3-dose series 4 weeks apart starting 6–12 months after successful transplant, regardless of Hib vaccination history

Hepatitis A vaccination

Routine vaccination

- **Not at risk but want protection from hepatitis A** (identification of risk factor not required): 2-dose series HepA (Havrix 6–12 months apart or Vaqta 6–18 months apart [minimum interval: 6 months]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 5 months])

Special situations

- **At risk for hepatitis A virus infection:** 2-dose series HepA or 3-dose series HepA-HepB as above
 - **Chronic liver disease** (e.g., persons with hepatitis B, hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice the upper limit of normal)
 - **HIV infection**
 - **Men who have sex with men**
 - **Injection or noninjection drug use**

- **Persons experiencing homelessness**
- **Work with hepatitis A virus** in research laboratory or with nonhuman primates with hepatitis A virus infection
- **Travel in countries with high or intermediate endemic hepatitis A** (HepA-HepB [Twinrix] may be administered on an accelerated schedule of 3 doses at 0, 7, and 21–30 days, followed by a booster dose at 12 months)
- **Close, personal contact with international adoptee** (e.g., household or regular babysitting) in first 60 days after arrival from country with high or intermediate endemic hepatitis A (administer dose 1 as soon as adoption is planned, at least 2 weeks before adoptee's arrival)
- **Pregnancy** if at risk for infection or severe outcome from infection during pregnancy
- **Settings for exposure, including** health care settings targeting services to injection or noninjection drug users or group homes and nonresidential day care facilities for developmentally disabled persons (individual risk factor screening not required)

Hepatitis B vaccination

Routine vaccination

- **Not at risk but want protection from hepatitis B** (identification of risk factor not required): 2- or 3-dose series (2-dose series Heplisav-B at least 4 weeks apart [2-dose series HepB only applies when 2 doses of Heplisav-B are used at least 4 weeks apart] or 3-dose series Engerix-B or Recombivax HB at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 8 weeks / dose 1 to dose 3: 16 weeks]) or 3-dose series HepA-HepB (Twinrix at 0, 1, 6 months [minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 5 months])

Special situations

- **At risk for hepatitis B virus infection:** 2-dose (Heplisav-B) or 3-dose (Engerix-B, Recombivax HB) series or 3-dose series HepA-HepB (Twinrix) as above
 - **Chronic liver disease** (e.g., persons with hepatitis C, cirrhosis, fatty liver disease, alcoholic liver disease, autoimmune hepatitis, alanine aminotransferase [ALT] or aspartate aminotransferase [AST] level greater than twice upper limit of normal)
 - **HIV infection**
 - **Sexual exposure risk** (e.g., sex partners of hepatitis B surface antigen [HBsAg]-positive persons; sexually active persons not in mutually monogamous relationships; persons seeking evaluation or treatment for a sexually transmitted infection; men who have sex with men)

- **Current or recent injection drug use**
- **Percutaneous or mucosal risk for exposure to blood** (e.g., household contacts of HBsAg-positive persons; residents and staff of facilities for developmentally disabled persons; health care and public safety personnel with reasonably anticipated risk for exposure to blood or blood-contaminated body fluids; hemodialysis, peritoneal dialysis, home dialysis, and predialysis patients; persons with diabetes mellitus age younger than 60 years, shared clinical decision-making for persons age 60 years or older)
- **Incarcerated persons**
- **Travel in countries with high or intermediate endemic hepatitis B**
- **Pregnancy** if at risk for infection or severe outcome from infection during pregnancy (Heplisav-B not currently recommended due to lack of safety data in pregnant women)

Human papillomavirus vaccination

Routine vaccination

- **HPV vaccination recommended for all persons through age 26 years:** 2- or 3-dose series depending on age at initial vaccination or condition:
 - **Age 15 years or older at initial vaccination:** 3-dose series at 0, 1–2 months, 6 months (minimum intervals: dose 1 to dose 2: 4 weeks / dose 2 to dose 3: 12 weeks / dose 1 to dose 3: 5 months; repeat dose if administered too soon)
 - **Age 9–14 years at initial vaccination and received 1 dose or 2 doses less than 5 months apart:** 1 additional dose
 - **Age 9–14 years at initial vaccination and received 2 doses at least 5 months apart:** HPV vaccination series complete, no additional dose needed
- **Interrupted schedules:** If vaccination schedule is interrupted, the series does not need to be restarted
- **No additional dose recommended after completing series with recommended dosing intervals using any HPV vaccine**

Shared clinical decision-making

- **Some adults age 27–45 years:** Based on shared clinical decision-making, 2- or 3-dose series as above

Special situations

- **Age ranges recommended above for routine and catch-up vaccination or shared clinical decision-making also apply in special situations**

Notes

Recommended Adult Immunization Schedule, United States, 2021

- **Immunocompromising conditions, including HIV infection:** 3-dose series as above, regardless of age at initial vaccination
- **Pregnancy:** HPV vaccination not recommended until after pregnancy; no intervention needed if vaccinated while pregnant; pregnancy testing not needed before vaccination

Influenza vaccination

Routine vaccination

- **Persons age 6 months or older:** 1 dose any influenza vaccine appropriate for age and health status annually
- For additional guidance, see www.cdc.gov/flu/professionals/index.htm

Special situations

- **Egg allergy, hives only:** 1 dose any influenza vaccine appropriate for age and health status annually
- **Egg allergy—any symptom other than hives** (e.g., angioedema, respiratory distress): 1 dose any influenza vaccine appropriate for age and health status annually. If using an influenza vaccine other than RIV4 or cclIV4, administer in medical setting under supervision of health care provider who can recognize and manage severe allergic reactions.
- Severe allergic reactions to any vaccine can occur even in the absence of a history of previous allergic reaction. Therefore, all vaccine providers should be familiar with the office emergency plan and certified in cardiopulmonary resuscitation.
- A previous severe allergic reaction to any influenza vaccine is a contraindication to future receipt of the vaccine.
- **LAIV4 should not be used** in persons with the following conditions or situations:
 - History of severe allergic reaction to any vaccine component (excluding egg) or to a previous dose of any influenza vaccine
 - Immunocompromised due to any cause (including medications and HIV infection)
 - Anatomic or functional asplenia
 - Close contacts or caregivers of severely immunosuppressed persons who require a protected environment
 - Pregnancy
 - Cranial CSF/oropharyngeal communications
 - Cochlear implant

- Received influenza antiviral medications oseltamivir or zanamivir within the previous 48 hours, peramivir within the previous 5 days, or baloxavir within the previous 17 days
- Adults 50 years or older
- **History of Guillain-Barré syndrome within 6 weeks after previous dose of influenza vaccine:** Generally, should not be vaccinated unless vaccination benefits outweigh risks for those at higher risk for severe complications from influenza

Measles, mumps, and rubella vaccination

Routine vaccination

- **No evidence of immunity to measles, mumps, or rubella:** 1 dose
 - **Evidence of immunity:** Born before 1957 (health care personnel, see below), documentation of receipt of MMR vaccine, laboratory evidence of immunity or disease (diagnosis of disease without laboratory confirmation is not evidence of immunity)

Special situations

- **Pregnancy with no evidence of immunity to rubella:** MMR contraindicated during pregnancy; after pregnancy (before discharge from health care facility), 1 dose
- **Nonpregnant women of childbearing age with no evidence of immunity to rubella:** 1 dose
- **HIV infection with CD4 count ≥ 200 cells/mm³ for at least 6 months and no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart; MMR contraindicated for HIV infection with CD4 count < 200 cells/mm³
- **Severe immunocompromising conditions:** MMR contraindicated
- **Students in postsecondary educational institutions, international travelers, and household or close, personal contacts of immunocompromised persons with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart if previously did not receive any doses of MMR or 1 dose if previously received 1 dose MMR
- **Health care personnel:**
 - **Born in 1957 or later with no evidence of immunity to measles, mumps, or rubella:** 2-dose series at least 4 weeks apart for measles or mumps or at least 1 dose for rubella

- **Born before 1957 with no evidence of immunity to measles, mumps, or rubella:** Consider 2-dose series at least 4 weeks apart for measles or mumps or 1 dose for rubella

Meningococcal vaccination

Special situations for MenACWY

- **Anatomical or functional asplenia (including sickle cell disease), HIV infection, persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use:** 2-dose series MenACWY-D (Menactra, Menveo or MenQuadfi) at least 8 weeks apart and revaccinate every 5 years if risk remains
- **Travel in countries with hyperendemic or epidemic meningococcal disease, microbiologists routinely exposed to *Neisseria meningitidis*:** 1 dose MenACWY (Menactra, Menveo or MenQuadfi) and revaccinate every 5 years if risk remains
- **First-year college students who live in residential housing (if not previously vaccinated at age 16 years or older) and military recruits:** 1 dose MenACWY (Menactra, Menveo or MenQuadfi)
- For MenACWY **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting (e.g., in community or organizational settings and among men who have sex with men) and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm

Shared clinical decision-making for MenB

- **Adolescents and young adults age 16–23 years (age 16–18 years preferred) not at increased risk for meningococcal disease:** Based on shared clinical decision-making, 2-dose series MenB-4C (Bexsero) at least 1 month apart or 2-dose series MenB-FHbp (Trumenba) at 0, 6 months (if dose 2 was administered less than 6 months after dose 1, administer dose 3 at least 4 months after dose 2); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series)

Special situations for MenB

- **Anatomical or functional asplenia (including sickle cell disease), persistent complement component deficiency, complement inhibitor (e.g., eculizumab, ravulizumab) use, microbiologists routinely exposed to *Neisseria meningitidis*:** 2-dose primary series MenB-4C (Bexsero) at least one month apart or

- MenB-4C (Bexsero) at least 1 month apart or 3-dose primary series MenB-FHbp (Trumenba) at 0, 1–2, 6 months (if dose 2 was administered at least 6 months after dose 1, dose 3 not needed); MenB-4C and MenB-FHbp are not interchangeable (use same product for all doses in series); 1 dose MenB booster 1 year after primary series and revaccinate every 2–3 years if risk remains
- **Pregnancy:** Delay MenB until after pregnancy unless at increased risk and vaccination benefits outweigh potential risks
- For MenB **booster dose recommendations** for groups listed under “Special situations” and in an outbreak setting (e.g., in community or organizational settings and among men who have sex with men) and additional meningococcal vaccination information, see www.cdc.gov/mmwr/volumes/69/rr/rr6909a1.htm

Pneumococcal vaccination

Routine vaccination

- **Age 65 years or older** (immunocompetent—see www.cdc.gov/mmwr/volumes/68/wr/mm6846a5.htm?s_cid=mm6846a5_w): 1 dose PPSV23
 - If PPSV23 was administered prior to age 65 years, administer 1 dose PPSV23 at least 5 years after previous dose

Shared clinical decision-making

- **Age 65 years or older** (immunocompetent): 1 dose PCV13 based on **shared clinical decision-making** if previously not administered.
 - PCV13 and PPSV23 should not be administered during the same visit
 - If both PCV13 and PPSV23 are to be administered, PCV13 should be administered first
 - PCV13 and PPSV23 should be administered at least 1 year apart

Special situations

- **Age 19–64 years with chronic medical conditions (chronic heart [excluding hypertension], lung, or liver disease, diabetes), alcoholism, or cigarette smoking:** 1 dose PPSV23

- **Age 19 years or older with immunocompromising conditions (congenital or acquired immunodeficiency [including B- and T-lymphocyte deficiency, complement deficiencies, phagocytic disorders, HIV infection], chronic renal failure, nephrotic syndrome, leukemia, lymphoma, Hodgkin disease, generalized malignancy, iatrogenic immunosuppression [e.g., drug or radiation therapy], solid organ transplant, multiple myeloma) or anatomical or functional asplenia (including sickle cell disease and other hemoglobinopathies):** 1 dose PCV13 followed by 1 dose PPSV23 at least 8 weeks later, then another dose PPSV23 at least 5 years after previous PPSV23; at age 65 years or older, administer 1 dose PPSV23 at least 5 years after most recent PPSV23 (note: only 1 dose PPSV23 recommended at age 65 years or older)
- **Age 19 years or older with cerebrospinal fluid leak or cochlear implant:** 1 dose PCV13 followed by 1 dose PPSV23 at least 8 weeks later; at age 65 years or older, administer another dose PPSV23 at least 5 years after PPSV23 (note: only 1 dose PPSV23 recommended at age 65 years or older)

Tetanus, diphtheria, and pertussis vaccination

Routine vaccination

- **Previously did not receive Tdap at or after age 11 years:** 1 dose Tdap, then Td or Tdap every 10 years

Special situations

- **Previously did not receive primary vaccination series for tetanus, diphtheria, or pertussis:** At least 1 dose Tdap followed by 1 dose Td or Tdap at least 4 weeks after Tdap and another dose Td or Tdap 6–12 months after last Td or Tdap (Tdap can be substituted for any Td dose, but preferred as first dose), Td or Tdap every 10 years thereafter
- **Pregnancy:** 1 dose Tdap during each pregnancy, preferably in early part of gestational weeks 27–36
- **Wound management:** Persons with 3 or more doses of tetanus-toxoid-containing vaccine: For clean and minor wounds, administer Tdap or Td if more than 10 years since last dose of tetanus-toxoid-containing vaccine; for all other wounds, administer Tdap or Td if more than 5 years since last dose of tetanus-toxoid-containing vaccine. Tdap is preferred for persons who have not previously received Tdap or whose Tdap history is unknown. If a tetanus-toxoid-containing vaccine is indicated for a pregnant woman, use Tdap. For detailed information, see www.cdc.gov/mmwr/volumes/69/wr/mm6903a5.htm

Varicella vaccination

Routine vaccination

- **No evidence of immunity to varicella:** 2-dose series 4–8 weeks apart if previously did not receive varicella-containing vaccine (VAR or MMRV [measles-mumps-rubella-varicella vaccine] for children); if previously received 1 dose varicella-containing vaccine, 1 dose at least 4 weeks after first dose
 - Evidence of immunity: U.S.-born before 1980 (except for pregnant women and health care personnel [see below]), documentation of 2 doses varicella-containing vaccine at least 4 weeks apart, diagnosis or verification of history of varicella or herpes zoster by a health care provider, laboratory evidence of immunity or disease

Special situations

- **Pregnancy with no evidence of immunity to varicella:** VAR contraindicated during pregnancy; after pregnancy (before discharge from health care facility), 1 dose if previously received 1 dose varicella-containing vaccine or dose 1 of 2-dose series (dose 2: 4–8 weeks later) if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **Health care personnel with no evidence of immunity to varicella:** 1 dose if previously received 1 dose varicella-containing vaccine; 2-dose series 4–8 weeks apart if previously did not receive any varicella-containing vaccine, regardless of whether U.S.-born before 1980
- **HIV infection with CD4 count ≥ 200 cells/mm³ with no evidence of immunity:** Vaccination may be considered (2 doses 3 months apart); VAR contraindicated for HIV infection with CD4 count < 200 cells/mm³
- **Severe immunocompromising conditions:** VAR contraindicated

Zoster vaccination


Routine vaccination

- **Age 50 years or older:** 2-dose series RZV (Shingrix) 2–6 months apart (minimum interval: 4 weeks; repeat dose if administered too soon), regardless of previous herpes zoster or history of zoster vaccine live (ZVL, Zostavax) vaccination (administer RZV at least 2 months after ZVL)

Special situations

- **Pregnancy:** Consider delaying RZV until after pregnancy if RZV is otherwise indicated.
- **Severe immunocompromising conditions (including HIV infection with CD4 count < 200 cells/mm³):** Recommended use of RZV under review

2021 Recommended Immunizations for Children from Birth Through 6 Years Old



Is your family growing? To protect your new baby against whooping cough, get a Tdap vaccine. The recommended time is the 27th through 36th week of pregnancy. Talk to your doctor for more details.

Shaded boxes indicate the vaccine can be given during shown age range.

NOTE:

If your child misses a shot, you don't need to start over. Just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.

FOOTNOTES:

- * Two doses given at least four weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.
 - § Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 months after the first dose. All children and adolescents over 24 months of age who have not been vaccinated should also receive 2 doses of HepA vaccine.
- If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he or she may need.*

See back page for more information on vaccine-preventable diseases and the vaccines that prevent them.



For more information, call toll-free
1-800-CDC-INFO (1-800-232-4636)
or visit
www.cdc.gov/vaccines/parents



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Disease	Vaccine	Disease spread by	Disease symptoms	Disease complications
Chickenpox	Varicella vaccine protects against chickenpox.	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs)
Diphtheria	DTaP* vaccine protects against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death
Hib	Hib vaccine protects against <i>Haemophilus influenzae</i> type b.	Air, direct contact	May be no symptoms unless bacteria enter the blood	Meningitis (infection of the covering around the brain and spinal cord), intellectual disability, epiglottitis (life-threatening infection that can block the windpipe and lead to serious breathing problems), pneumonia (infection in the lungs), death
Hepatitis A	HepA vaccine protects against hepatitis A.	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders
Hepatitis B	HepB vaccine protects against hepatitis B.	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer
Influenza (Flu)	Flu vaccine protects against influenza.	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs)
Measles	MMR** vaccine protects against measles.	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death
Mumps	MMR** vaccine protects against mumps.	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness
Pertussis	DTaP* vaccine protects against pertussis (whooping cough).	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death
Polio	IPV vaccine protects against polio.	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death
Pneumococcal	PCV13 vaccine protects against pneumococcus.	Air, direct contact	May be no symptoms, pneumonia (infection in the lungs)	Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death
Rotavirus	RV vaccine protects against rotavirus.	Through the mouth	Diarrhea, fever, vomiting	Severe diarrhea, dehydration
Rubella	MMR** vaccine protects against rubella.	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects
Tetanus	DTaP* vaccine protects against tetanus.	Exposure through cuts in skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death

* DTaP combines protection against diphtheria, tetanus, and pertussis.

** MMR combines protection against measles, mumps, and rubella.

Talk to your child’s doctor or nurse about the vaccines recommended for their age. COVID-19 vaccination is recommended for some adolescents.

	Flu Influenza	Tdap Tetanus, diphtheria, pertussis	HPV Human papillomavirus	Meningococcal		Pneumococcal	Hepatitis B	Hepatitis A	Polio	MMR Measles, mumps, rubella	Chickenpox Varicella
				MenACWY	MenB						
7-8 Years	Green	Orange		Purple		Purple	Orange	Orange	Orange	Orange	Orange
9-10 Years	Green	Orange	Purple, Blue	Purple		Purple	Orange	Orange	Orange	Orange	Orange
11-12 Years	Green	Green, Orange	Green	Green, Orange		Purple	Orange	Orange	Orange	Orange	Orange
13-15 Years	Green	Orange	Orange	Orange		Purple	Orange	Orange	Orange	Orange	Orange
16-18 Years	Green	Orange	Orange	Green, Orange		Purple	Orange	Orange	Orange	Orange	Orange

More information:


Everyone 6 months and older should get a flu vaccine every year.


All 11- through 12-year olds should get one shot of Tdap.


All 11- through 12-year olds should get a 2-shot series of HPV vaccine. A 3-shot series is needed for those with weakened immune systems and those who start the series at 15 years or older.


All 11- through 12-year olds should get one shot of meningococcal conjugate (MenACWY). A booster shot is recommended at age 16.

Teens 16–18 years old **may** be vaccinated with a serogroup B meningococcal (MenB) vaccine.

 These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

 These shaded boxes indicate the vaccine should be given if a child is catching up on missed vaccines.

 These shaded boxes indicate the vaccine is recommended for children with certain health or lifestyle conditions that put them at an increased risk for serious diseases. See vaccine-specific recommendations at www.cdc.gov/vaccines/hcp/acip-recs/.

 This shaded box indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™



Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Disease	Vaccine	Disease spread by	Disease symptoms	Disease complications
Chickenpox	Varicella vaccine protects against chickenpox.	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs)
Diphtheria	Tdap* and Td** vaccines protect against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death
Hepatitis A	HepA vaccine protects against hepatitis A.	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders
Hepatitis B	HepB vaccine protects against hepatitis B.	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer
Human Papillomavirus	HPV vaccine protects against human papillomavirus.	Direct skin contact	May be no symptoms, genital warts	Cervical, vaginal, vulvar, penile, anal, oropharyngeal cancers
Influenza (Flu)	Flu vaccine protects against influenza.	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs)
Measles	MMR*** vaccine protects against measles.	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death
Meningococcal Disease	MenACWY and MenB vaccines protect against meningococcal disease.	Air, direct contact	Sudden onset of fever, headache, and stiff neck, dark purple rash	Loss of limb, deafness, nervous system disorders, developmental disabilities, seizure disorder, stroke, death
Mumps	MMR*** vaccine protects against mumps.	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness
Pertussis	Tdap* vaccine protects against pertussis.	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death
Pneumococcal Disease	Pneumococcal vaccine protects against pneumococcal disease.	Air, direct contact	May be no symptoms, pneumonia (infection in the lungs)	Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death
Polio	Polio vaccine protects against polio.	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death
Rubella	MMR*** vaccine protects against rubella.	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects
Tetanus	Tdap* and Td** vaccines protect against tetanus.	Exposure through cuts on skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death

*Tdap combines protection against diphtheria, tetanus, and pertussis.

**Td combines protection against diphtheria and tetanus.

***MMR combines protection against measles, mumps, and rubella.

If you have any questions about your child's vaccines, talk to your child's doctor or nurse.

COVID-19 Vaccine

Quick Reference Guide for
Healthcare Professionals



The table below provides basic information on the proper storage, preparation, and administration of the currently authorized COVID-19 vaccine products in the United States. For additional information and detailed clinical guidance go to the manufacturer's and CDC's webpages listed.

		Pfizer	Moderna	Janssen
GENERAL	EUA	www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/pfizer-biontech-covid-19-vaccine	www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/moderna-covid-19-vaccine	www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/janssen-covid-19-vaccine
	CDC Vaccine Information	www.cdc.gov/vaccines/covid-19/info-by-product/pfizer/index.html	www.cdc.gov/vaccines/covid-19/info-by-product/moderna/index.html	www.cdc.gov/vaccines/covid-19/info-by-product/janssen/index.html
	Manufacturer Contact information	Website: www.cvdvaccine.com Medical information: 800-438-1985 Customer service: 800-879-3477	Website: www.modernatx.com Medical Information: 866-663-3762	Website: www.vaxcheck.inj Medical information: 1-800-565-4008
STORAGE & HANDLING	How supplied	Multidose vial: 6 doses	Multidose vial: Maximum of 11 doses Multidose vial: Maximum of 15 doses	Multidose vial: 5 doses
	Diluent	0.9% sodium chloride (preservative-free, normal saline) provided in the ancillary kit. Do NOT use other diluent.	None	None
	Storage Temperatures: Before Puncture	Between: -80°C and -60°C (-112°F and -76°F) until the expiration date -25°C and -15°C (-13°F and 5°F) for up to 2 weeks 2°C and 8°C (36°F and 46°F) for up to 1 month (31 days).	Between: -50°C and -15°C (-58°F and 5°F) until the expiration date 2°C and 8°C (36°F and 46°F) for up to 30 days 8°C and 25°C (46° and 77°F) for a total of 24 hours	Between: 2°C and 8°C (36°F and 46°F) until the expiration date.
	Storage Temperatures: After puncture	Between: 2°C to 25°C (36°F to 77°F) for up to 6 hours. Discard any unused vaccine after 6 hours.	Between: 2°C and 25°C (36°F and 77°F) for up to 12 hours. Discard any unused vaccine after 12 hours.	Between: 2°C and 8°C (36°F and 46°F) for up to 6 hours. 9°C and 25°C (47°F and 77°F) for up to 2 hours. Discard any unused vaccine after these time frames.
	Transport Temperatures: Before Puncture	Between: -80°C and -60°C (-112°F and -76°F) -25°C and -15°C (-13°F and 5°F) 2°C and 8°C (36°F and 46°F)	Between: -50°C and -15°C (-58°F and 5°F) 2°C and 8°C (36°F and 46°F) for up to 12 cumulative hours.	Between: 2°C and 8°C (36°F and 46°F)
	Transport Temperatures*: After Puncture	Between: 2°C to 25°C (36°F to 77°F) for up to 6 hours.	Between: 2°C and 25°C (36°F and 77°F) for up to 12 hours.	Between: 2°C and 8°C (36°F and 46°F) for up to 6 hours
Type of Vaccine	mRNA	mRNA	Viral vector	
Age Indications	12 years of age and older	18 years of age and older	18 years of age and older	
Schedule[†]	2-doses, separated by 21 days. Both doses must be Pfizer-BioNTech vaccine	2 doses, separated by 28 days. Both doses should be Moderna vaccine	1 dose only	
Dosage	0.3 mL	0.5 mL	0.5 mL	
Needle gauge/length	12 through 18 years of age: 22–25 gauge, 1" 19 years of age and older: 22–25 gauge, 1 – 1½"	22–25 gauge, 1 – 1½"	22–25 gauge, 1 – 1½"	

*CDC recommends transporting vaccine at refrigerated or frozen temperatures.

[†]COVID-19 vaccines and other vaccines may be administered on the same day, as well as within 14 days of each other. When deciding if to administer COVID-19 vaccines and other vaccines, providers should consider whether the patient is behind or at risk of becoming behind on recommended vaccines, their risk of vaccine-preventable diseases (e.g., during an outbreak), and the reactogenicity profile of the vaccines.

COVID-19 Vaccine

Quick Reference Guide for
Healthcare Professionals



	Pfizer	Moderna	Janssen
Route	Intramuscular (IM) injection	Intramuscular (IM) injection	Intramuscular (IM) injection
Site	Deltoid	Deltoid	Deltoid
Thawing Frozen Vaccine	Between: 2°C and 8°C (36°F and 46°F) or Room temperature up to 25°C (77°F) Do NOT refreeze thawed vaccine.	Between: 2°C and 8°C (36°F and 46°F) or 8°C to 25°C (46°F to 77°F) Do NOT refreeze thawed vaccine.	N/A
Mixing Vaccine	Mix vaccine with 1.8 mL of 0.9% sodium chloride (preservative-free, normal saline)	Do NOT mix with any diluent	Do NOT mix with any diluent
VACCINE ADMINISTRATION	<p>Contraindications</p> <ul style="list-style-type: none"> Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the COVID-19 vaccine Immediate allergic reaction† of any severity to a previous dose or known (diagnosed) allergy to a component of the vaccine <p>Note: Persons who have a contraindication to an mRNA COVID-19 vaccine (Moderna or Pfizer-BioNTech) may be able to receive the Janssen COVID-19 vaccine (see footnote).‡</p> <p>Persons who have a contraindication to Janssen COVID-19 vaccine may be able to receive an mRNA COVID-19 vaccine (see footnote).‡</p> <p>Contraindications/Precautions</p> <p>Precautions</p> <ul style="list-style-type: none"> Most people determined to have a precaution to a COVID-19 vaccine at their appointment can and should be administered vaccine. History of an immediate allergic reaction† to any other vaccine or injectable therapy (i.e., intramuscular, intravenous, or subcutaneous vaccines or therapies) <ul style="list-style-type: none"> » This includes people with a reaction to a vaccine or injectable therapy that contains multiple components, one of which is a vaccine component, but for whom it is unknown which component elicited the immediate allergic reaction. People with a contraindication to mRNA COVID-19 vaccines have a precaution to Janssen COVID-19 Vaccine, and vice versa. (see footnote).‡ Moderate to severe acute illness <p>See Interim Clinical Considerations for Use of mRNA COVID-19 Vaccines Currently Authorized in the United States www.cdc.gov/vaccines/covid-19/info-by-product/clinical-considerations.html</p>		
	Post-Vaccination Observation	<p>30 minutes: Persons with a history of an immediate allergic reaction of any severity to any other vaccine or injectable therapy or a history of anaphylaxis (from any cause)</p> <p>15 minutes: All other persons</p>	
Most common adverse events	<p>Injection site: pain, swelling, redness</p> <p>Systemic: fatigue, headache, muscle pain, chills, fever, joint pain</p>	<p>Injection site: pain, swelling, redness</p> <p>Systemic: fatigue, headache, muscle pain, chills, fever, nausea, joint pain</p>	<p>Injection site: pain, redness, swelling</p> <p>Systemic: fatigue, headache, muscle pain, nausea, fever</p>

†For the purpose of this guidance, an immediate allergic reaction is defined as any hypersensitivity-related signs or symptoms, such as urticaria, angioedema, respiratory distress (e.g., wheezing, stridor), or anaphylaxis that occur within 4 hours following exposure to a vaccine or medication.

‡Consider consultation with an allergist-immunologist to help determine if the patient can safely receive vaccination. Healthcare providers and health departments may also request a consultation from the Clinical Immunization Safety Assessment COVID vax Project <https://www.cdc.gov/vaccinesafety/ensuringsafety/monitoring/cisa/index.html>. Vaccination of these individuals should only be done in an appropriate setting under the supervision of a healthcare provider experienced in the management of severe allergic reactions.

- People with a contraindication to mRNA COVID-19 vaccines (including due to a known PEG allergy) have a precaution to Janssen COVID-19 vaccination. People who have previously received an mRNA COVID-19 vaccine dose but have a contraindication to a second dose should wait at least 28 days to receive Janssen COVID-19 vaccine.
- People with a contraindication to Janssen COVID-19 vaccine (including due to a known polysorbate allergy) have a precaution to mRNA COVID-19 vaccination.