

Prior Authorization Requirements

Elevate Medicare Choice (HMO D-SNP) and Elevate Medicare Select (HMO)

Effective: 05/01/2025

ABALOPARATIDE

Products Affected

• TYMLOS

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 MONTHS |
| Other Criteria | OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ABATACEPT IV

Products Affected

• ORENCIA (WITH MALTOSE)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. |
| Coverage Duration | RA, PJIA, PSA: INITIAL: 6 MOS, RENEWAL: 12 MOS. ACUTE GRAFT VERSUS HOST DISEASE (AGVHD): 1 MO. |
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ABATACEPT SQ

- ORENCIA
- ORENCIA CLICKJECT

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ABEMACICLIB

Products Affected

VERZENIO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ABIRATERONE

- abiraterone
- abirtega

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC), METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ABIRATERONE SUBMICRONIZED

Products Affected

YONSA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ACALABRUTINIB

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | PREVIOUSLY TREATED MANTLE CELL LYMPHOMA: INTOLERANCE TO BRUKINSA. CHRONIC LYMPHOCYTIC LEUKEMIA OR SMALL LYMPHOCYTIC LYMPHOMA: INTOLERANCE TO BRUKINSA OR IMBRUVICA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ADAGRASIB

Products Affected

KRAZATI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ADALIMUMAB

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)

- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST |
| Coverage Duration | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------------------|---|
| PA Criteria Other Criteria | Criteria Details INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR |
| | USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: NO ISOLATED ANTERIOR |
| | UVEITIS. RENEWAL: RA, HS, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |

| PA Criteria | Criteria Details |
|------------------------|---|
| | MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ADALIMUMAB-AATY

- YUFLYMA(CF)
- YUFLYMA(CF) AI CROHN'S-UC-HS
- YUFLYMA(CF) AUTOINJECTOR

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST |
| Coverage Duration | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|--|
| Other Criteria | INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, HS, UVEITIS: CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER |
| | SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR |

| PA Criteria | Criteria Details |
|------------------------|---|
| | PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ADALIMUMAB-ADBM

- CYLTEZO(CF)
- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS
- CYLTEZO(CF) PEN PSORIASIS-UV

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST |
| Coverage Duration | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------------------|--|
| PA Criteria Other Criteria | INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT |
| | USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |
| | MOLECULES FOR HS. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, HS, UVEITIS: CONTINUES TO BENEFIT FROM MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM |
| | MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR |

| PA Criteria | Criteria Details |
|------------------------|---|
| | PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AFATINIB

Products Affected

• GILOTRIF

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ALECTINIB

Products Affected

ALECENSA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ALPELISIB-PIQRAY

Products Affected

 PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AMIKACIN LIPOSOMAL INH

Products Affected

ARIKAYCE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | MYCOBACTERIUM AVIUM COMPLEX (MAC) LUNG DISEASE: RENEWAL: 1) NO POSITIVE MAC SPUTUM CULTURE AFTER CONSECUTIVE NEGATIVE CULTURES, AND 2) IMPROVEMENT IN SYMPTOMS. ADDITIONALLY, FOR FIRST RENEWAL, APPROVAL REQUIRES AT LEAST ONE NEGATIVE SPUTUM CULTURE FOR MAC BY SIX MONTHS OF ARIKAYCE TREATMENT. FOR SECOND AND SUBSEQUENT RENEWALS, APPROVAL REQUIRES AT LEAST THREE NEGATIVE SPUTUM CULTURES FOR MAC BY 12 MONTHS OF ARIKAYCE TREATMENT. |
| Age Restrictions | |
| Prescriber Restrictions | MAC LUNG DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR INFECTIOUS DISEASE SPECIALIST. |
| Coverage Duration | INITIAL/RENEWAL: 6 MONTHS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AMIVANTAMAB-VMJW

Products Affected

RYBREVANT

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ANAKINRA

Products Affected

KINERET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS. |
| Required Medical Information | INITIAL: CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR \$100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES. DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. |
| Coverage Duration | RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. CAPS, DIRA: LIFETIME. |
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. CAPS, DIRA: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Off Label Uses | |
| Part B Prerequisite | No |

APALUTAMIDE

Products Affected

• ERLEADA ORAL TABLET 240 MG, 60 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

APOMORPHINE - ONAPGO

Products Affected

ONAPGO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | PD: INITIAL: 1) RESPONSIVE TO LEVODOPA, 2) CURRENT MEDICATION REGIMEN, INCLUDING LEVODOPA, HAS BEEN AT A STABLE DOSE FOR AT LEAST 28 DAYS, 3) MOTOR SYMPTOMS ARE CURRENTLY UNCONTROLLED (DEFINED AS AN AVERAGE OFF TIME OF AT LEAST 3 HOURS/DAY, FOR AT LEAST 2 HOURS EACH DAY), AND 4) DOES NOT HAVE ANY OF THE FOLLOWING: ORTHOSTATIC HYPOTENSION, HISTORY OF PROLONGED QTC (GREATER THAN 450 MSEC FOR MALE OR GREATER THAN 470 MSEC FOR FEMALE), ACTIVE OR UNCONTROLLED PSYCHOSIS, ACTIVE OR UNCONTROLLED DEPRESSION. RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

APOMORPHINE - SL

Products Affected

 KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER. |
| Prescriber Restrictions | PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | PD: RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF THERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

APREMILAST

- OTEZLA
- OTEZLA STARTER

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: MILD PLAQUE PSORIASIS (PSO): 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC THERAPY (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) OR ONE CONVENTIONAL TOPICAL THERAPY (E.G., PUVA [PHOTOTHERAPY], UVB [ULTRAVIOLET LIGHT B], TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR MODERATE TO SEVERE PSO. BEHCETS DISEASE: 1) HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: MILD PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION, PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. MODERATE TO SEVERE PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. MODERATE TO SEVERE PSO: 1) CONTUNUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. MODERATE TO SEVERE PSO: 1) CONTUNUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR MODERATE TO SEVERE PSO: 1) |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ARIMOCLOMOL

Products Affected

MIPLYFFA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | NIEMANN-PICK DISEASE TYPE C (NPC): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEUROLOGIST OR GENETICIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | NPC: RENEWAL: IMPROVEMENT OR SLOWING OF DISEASE PROGRESSION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ASCIMINIB

Products Affected

• SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PREVIOUSLY TREATED OR T315I MUTATION PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ASFOTASE ALFA

Products Affected

• STRENSIQ

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|---|
| Other Criteria | INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6 MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II) CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V) NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM CALCIUM, (VI) HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC DEFORMITIES, (II) PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OR PRESENCE OF NONTRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) |
| | INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

ATOGEPANT

Products Affected

• QULIPTA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AVACOPAN

Products Affected

TAVNEOS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | ANTI-NEUTROPHIL CYTOPLASMIC AUTOANTIBODY (ANCA)-ASSOCIATED VASCULITIS: INITIAL: ANCA SEROPOSITIVE (ANTI-PR3 OR ANTI-MPO). |
| Age Restrictions | |
| Prescriber Restrictions | ANCA-ASSOCIATED VASCULITIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 6 MONTHS. |
| Other Criteria | ANCA-ASSOCIATED VASCULITIS: RENEWAL: CONTINUES TO BENEFIT FROM THERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AVAPRITINIB

Products Affected

AYVAKIT

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AXATILIMAB-CSFR

Products Affected

NIKTIMVO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AXITINIB

Products Affected

• INLYTA ORAL TABLET 1 MG, 5 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AZACITIDINE

Products Affected

ONUREG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

AZTREONAM INHALED

Products Affected

CAYSTON

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | 7 YEARS OF AGE OR OLDER |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BEDAQUILINE

Products Affected

• SIRTURO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 WEEKS |
| Other Criteria | PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BELIMUMAB

Products Affected

• BENLYSTA SUBCUTANEOUS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BELUMOSUDIL

Products Affected

REZUROCK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BELZUTIFAN

Products Affected

WELIREG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BENDAMUSTINE

Products Affected

- bendamustine intravenous recon soln
- BENDAMUSTINE INTRAVENOUS SOLUTION
- BENDEKA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BENRALIZUMAB

Products Affected

- FASENRA
- FASENRA PEN

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. |
| Age Restrictions | |
| Prescriber Restrictions | ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS (EGPA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-2 INHIBITOR) FOR EGPA. RENEWAL: ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. EGPA: 1) REDUCTION IN GPA SYMPTOMS COMPARED TO BASELINE OR ABILITY TO REDUCE/ELIMINATE CORTICOSTEROID USE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR EGPA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BETAINE

Products Affected

• betaine

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BEVACIZUMAB-ADCD

Products Affected

VEGZELMA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BEVACIZUMAB-AWWB

Products Affected

• MVASI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BEVACIZUMAB-BVZR

Products Affected

ZIRABEV

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BEXAROTENE

Products Affected

• bexarotene

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BINIMETINIB

Products Affected

MEKTOVI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BORTEZOMIB

Products Affected

- bortezomib injectionBORUZU

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BOSENTAN

Products Affected

• bosentan

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE. RENEWAL: NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BOSUTINIB

Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

BRIGATINIB

Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS, DOSE PACK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

C1 ESTERASE INHIBITOR-HAEGARDA

Products Affected

• HAEGARDA SUBCUTANEOUS RECON SOLN 2,000 UNIT, 3,000 UNIT

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING COMPLEMENT TESTING: C1INH PROTEIN LEVELS, C4 PROTEIN LEVELS, C1-INH FUNCTIONAL LEVELS, C1Q. |
| Age Restrictions | |
| Prescriber Restrictions | HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, ALLERGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CABOZANTINIB CAPSULE

Products Affected

 COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CABOZANTINIB TABLET

Products Affected

• CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CANNABIDIOL

Products Affected

• EPIDIOLEX

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CAPIVASERTIB

Products Affected

• TRUQAP

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CAPMATINIB

Products Affected

TABRECTA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CARGLUMIC ACID

Products Affected

• carglumic acid

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: ACUTE OR CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS. |
| Other Criteria | RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CERITINIB

Products Affected

· ZYKADIA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CERTOLIZUMAB PEGOL

Products Affected

- CIMZIA POWDER FOR RECONST
- CIMZIA SUBCUTANEOUS SYRINGE KIT 400 MG/2 ML (200 MG/ML X 2)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI). |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, SKYRIZI, TREMFYA, OTEZLA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, CD: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: STELARA/SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. INITIAL FOR RA, PSA, PSO, AS, CD, PJIA: TRIAL OF OR CONTRAINDICATION TO THE STEP AGENTS IS NOT REQUIRED IF THE PATIENT IS PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT. INITIAL/RENEWAL FOR PSA, PSO, AS, CD, NR-AXSPA, PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR SAME INDICATION. RENEWAL FOR RA, PSA, AS, PSO, NR-AXSPA, PJIA: CONTINUES TO BENEFIT FROM MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CETUXIMAB

Products Affected

• ERBITUX

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CLADRIBINE

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 48 WEEKS. |
| Other Criteria | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CLOBAZAM-SYMPAZAN

Products Affected

SYMPAZAN

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | LGS: 1) UNABLE TO TAKE TABLETS OR SUSPENSIONS, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF CLOBAZAM. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

COBIMETINIB

Products Affected

COTELLIC

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CORTICOTROPIN

- ACTHAR
- ACTHAR SELFJECT SUBCUTANEOUS PEN INJECTOR 40 UNIT/0.5 ML, 80 UNIT/ML

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES. |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST. |
| Coverage Duration | INFANTILE SPASMS AND MS: 28 DAYS. ALL OTHER FDA APPROVED INDICATIONS: INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | Yes |

CRIZOTINIB CAPSULE

Products Affected

• XALKORI ORAL CAPSULE

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

CRIZOTINIB PELLETS

Products Affected

• XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | NON-SMALL CELL LUNG CANCER (NSCLC), ANAPLASTIC LARGE CELL LYMPHOMA (ALCL), INFLAMMATORY MYOFIBROBLASTIC TUMOR (IMT): UNABLE TO SWALLOW CAPSULES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DABRAFENIB CAPSULES

Products Affected

• TAFINLAR ORAL CAPSULE

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DABRAFENIB SUSPENSION

Products Affected

 TAFINLAR ORAL TABLET FOR SUSPENSION

| PA Criteria | Criteria Details |
|------------------------------------|---------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | UNABLE TO SWALLOW TAFINILAR CAPSULES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DACOMITINIB

Products Affected

VIZIMPRO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DALFAMPRIDINE

Products Affected

• dalfampridine

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DAROLUTAMIDE

Products Affected

NUBEQA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS |
| Other Criteria | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MHSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DASATINIB

Products Affected

• dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND DASATINIB IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DATOPOTAMAB DERUXTECAN-DLNK

Products Affected

DATROWAY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DECITABINE/CEDAZURIDINE

Products Affected

• INQOVI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DEFERASIROX

- deferasirox oral granules in packetdeferasirox oral tablet

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). CHRONIC IRON OVERLOAD IN NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS), AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G OF DRY LIVER WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G OF DRY LIVER WEIGHT OR GREATER. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL (CHRONIC IRON OVERLOAD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL (CHRONIC IRON OVERLOAD): DEFERASIROX SPRINKLE PACKETS: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX ORAL TABLET OR TABLET FOR ORAL SUSPENSION. |
| Indications | All FDA-approved Indications. |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Off Label Uses | |
| Part B Prerequisite | No |

DENOSUMAB-XGEVA

Products Affected

• XGEVA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DEUTETRABENAZINE

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 18
- MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4)

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | TARDIVE DYSKINESIA: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DICLOFENAC TOPICAL SOLUTION

Products Affected

• diclofenac sodium topical solution in metered-dose pump

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 MONTHS |
| Other Criteria | OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DIMETHYL FUMARATE

Products Affected

• dimethyl fumarate oral capsule,delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DIROXIMEL FUMARATE

Products Affected

VUMERITY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DOSTARLIMAB-GXLY

Products Affected

JEMPERLI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DRONABINOL CAPSULE

Products Affected

• dronabinol

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 MONTHS |
| Other Criteria | NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DROXIDOPA

Products Affected

droxidopa

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION. |
| Age Restrictions | |
| Prescriber Restrictions | NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST. |
| Coverage Duration | INITIAL: 3 MONTHS RENEWAL: 12 MONTHS |
| Other Criteria | NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DUPILUMAB

- DUPIXENT PEN
- DUPIXENT SYRINGE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL OF 150 TO 1500 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY. ATOPIC DERMATITIS (AD): AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: AD, PRURIGO NODULARIS (PN): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST. EOSINOPHILIC COPD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. |
| Coverage Duration | INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA, COPD: 12 MOS. RENEWAL: ALL INDICATIONS: 12 MOS. |

| Other Criteria INITIAL: AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
|---|----------------|--|
| CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | PA Criteria | Criteria Details |
| OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | Other Criteria | |
| (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| INHIBITOR, OR JAK INHIBITOR), AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | |
| OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING |
| AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, |
| WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA | | OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY |
| | | |
| CVMDTOMO MODE THAN TWICE/WILLIAM AND AND MICHT | | |
| | | SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT |
| WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR | | |
| SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY | | |
| LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE | | |
| WITH XOLAIR, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS | | |
| WHEN USED FOR ASTHMA. CHRONIC RHINOSINUSITIS WITH | | |
| NASAL POLYPS (CRSWNP): 1) A 56 DAY TRIAL OF ONE TOPICAL | | |
| NASAL CORTICOSTEROID, AND 2) NO CONCURRENT USE WITH | | |
| ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL | | |
| MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN | | |
| AUTOIMMUNE INDICATION. PN: 1) CHRONIC PRURITIS (ITCH | | |
| MORE THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, | | |
| AND HISTORY OR SIGN OF A PROLONGED SCRATCHING | | |
| BEHAVIOR, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE | | |
| TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL). EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A | | |
| LAMA/LABA/ICS, AND 2) NO CONCURRENT USE WITH | | |
| ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL | | |
| ANOTHER STSTEMIC BIOLOGIC OR TAROLTED SWALL | | ANOTHER STSTEMIC DIOLOGIC OR TARGETED SWILL |

| PA Criteria | Criteria Details |
|------------------------|---|
| | MOLECULES FOR EOSINOPHILIC COPD. RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. EOE: IMPROVEMENT WHILE ON THERAPY. CRSWNP: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AN AUTOIMMUNE INDICATION. ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITIS OR PRURIGINOUS LESIONS. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR EOSINOPHILIC COPD, AND 3) CLINICAL RESPONSE AS EVIDENCED BY (A) REDUCTION IN COPD EXACERBATIONS FROM BASELINE, (B) REDUCTION IN SEVERITY OR FREQUENCY OF COPD-RELATED SYMPTOMS, OR (C) INCREASE IN FEV1 OF AT LEAST 5 PERCENT FROM PRETREATMENT BASELINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

DUVELISIB

Products Affected

COPIKTRA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

EFLORNITHINE

Products Affected

• IWILFIN

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ELACESTRANT

Products Affected

• ORSERDU ORAL TABLET 345 MG, 86 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ELAGOLIX

Products Affected

• ORILISSA ORAL TABLET 150 MG, 200 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. |
| Age Restrictions | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER. |
| Prescriber Restrictions | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS |
| Other Criteria | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND A PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ELRANATAMAB-BCMM

- ELREXFIO 44 MG/1.1 ML VIAL INNER, SUV, P/F
- ELREXFIO SUBCUTANEOUS SOLUTION 40 MG/ML

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ELTROMBOPAG - ALVAIZ

Products Affected

ALVAIZ

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT IS LESS THAN 30 X 10^9/L FROM AT LEAST 2 SEPARATE LABS IN THE LAST 3 MONTHS, OR 2) PLATELET COUNT IS LESS THAN 50 X 10^9/L FROM AT LEAST 2 SEPARATE LABS IN THE LAST 3 MONTHS AND HAD A PRIOR BLEEDING EVENT. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST. |
| Coverage Duration | ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO. |
| Other Criteria | INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS) OR SPLEEN TYROSINE KINASE (SYK) INHIBITOR. RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNT FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS OR SYK INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ELTROMBOPAG - PROMACTA

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT OF LESS THAN 30 X 10^9/L FROM AT LEAST 2 SEPARATE LAB TESTS IN THE LAST 3 MONTHS, OR 2) PLATELET COUNT OF LESS THAN 50 X 10^9/L FROM AT LEAST 2 SEPARATE LAB TESTS IN THE LAST 3 MONTHS AND A PRIOR BLEEDING EVENT. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST. |
| Coverage Duration | ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO. |
| Other Criteria | INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR HAD AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS) OR SPLEEN TYROSINE KINASE (SYK) INHIBITOR. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLET OR PATIENT IS UNABLE TOLERATE TABLET FORMULATION. RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNTS FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS OR SYK INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ENASIDENIB

Products Affected

• IDHIFA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ENCORAFENIB

Products Affected

BRAFTOVI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ENTRECTINIB CAPSULES

Products Affected

 ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ENTRECTINIB PELLETS

Products Affected

 ROZLYTREK ORAL PELLETS IN PACKET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), SOLID TUMORS: 1) TRIAL OF OR CONTRAINDICATION TO ROZLYTREK CAPSULES MADE INTO AN ORAL SUSPENSION, AND 2) DIFFICULTY OR UNABLE TO SWALLOW CAPSULES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ENZALUTAMIDE

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: ALL INDICATIONS: 12 MONTHS. RENEWAL: MCRPC, NMCRPC, MCSPC: 12 MONTHS. |
| Other Criteria | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NON-METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (NMCSPC): HIGH RISK FOR METASTASIS (I.E. PSA DOUBLING TIME OF 9 MONTHS OR LESS). METASTATIC CRPC (MCRPC), NMCRPC, METASTATIC CSPC (MCSPC), NMCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: MCRPC, NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

EPCORITAMAB-BYSP

Products Affected

EPKINLY

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

EPOETIN ALFA-EPBX

Products Affected

 RETACRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL IS LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL IS LESS THAN 13G/DL. RENEWAL: 1) CKD IN ADULTS NOT ON DIALYSIS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 2) CKD IN PEDIATRIC PATIENTS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS APPROACHED OR EXCEEDS 12G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 3) ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. 4) CANCER CHEMOTHERAPY: (A) HEMOGLOBIN LEVEL IS LESS THAN 10 G/DL, OR (B) HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH. |
| Other Criteria | RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ERDAFITINIB

Products Affected

• BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ERLOTINIB

Products Affected

• erlotinib oral tablet 100 mg, 150 mg, 25 mg

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ESKETAMINE

Products Affected

SPRAVATO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST. |
| Coverage Duration | INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS. |
| Other Criteria | INITIAL: TRD, MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ETANERCEPT

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHE |
| | CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

EVEROLIMUS-AFINITOR

- everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg
 torpenz oral tablet 10 mg, 2.5 mg, 5 mg, 7.5
- mg

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

EVEROLIMUS-AFINITOR DISPERZ

Products Affected

• everolimus (antineoplastic) oral tablet for suspension

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FECAL MICROBIOTA CAPSULE

Products Affected

VOWST

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 30 DAYS |
| Other Criteria | CLOSTRIDIOIDES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FEDRATINIB

Products Affected

INREBIC

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FENFLURAMINE

Products Affected

FINTEPLA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS. |
| Other Criteria | INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FENTANYL CITRATE

Products Affected

• fentanyl citrate buccal lozenge on a handle

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FEZOLINETANT

Products Affected

VEOZAH

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | MENOPAUSAL VASOMOTOR SYMPTOMS (VMS): INITIAL: 1) EXPERIENCES 7 OR MORE HOT FLASHES PER DAY, AND 2) TRIAL OF OR CONTRAINDICATION TO HORMONAL THERAPY (E.G., ESTRADIOL TRANSDERMAL PATCH, ORAL CONJUGATED ESTROGENS). RENEWAL: 1) CONTINUED NEED FOR VMS TREATMENT (I.E., PERSISTENT HOT FLASHES), AND 2) REDUCTION IN VMS FREQUENCY OR SEVERITY DUE TO VEOZAH TREATMENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FILGRASTIM-AAFI

Products Affected

NIVESTYM

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FINERENONE

Products Affected

KERENDIA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FINGOLIMOD

Products Affected

• fingolimod

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FOSCARBIDOPA-FOSLEVODOPA

Products Affected

VYALEV

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | PD: INITIAL: 1) RESPONSIVE TO LEVODOPA, 2) CURRENT REGIMEN INCLUDES AT LEAST 400 MG/DAY OF LEVODOPA, AND 3) MOTOR SYMPTOMS ARE CURRENTLY UNCONTROLLED (DEFINED AS AN AVERAGE OFF TIME OF AT LEAST 2.5 HOURS/DAY OVER 3 CONSECUTIVE DAYS WITH A MINIMUM OF 2 HOURS EACH DAY). RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FREMANEZUMAB-VFRM

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. |
| Other Criteria | MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FRUQUINTINIB

Products Affected

• FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

FUTIBATINIB

Products Affected

 LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GALCANEZUMAB-GNLM

- EMGALITY PEN
- EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS. |
| Other Criteria | INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GANAXOLONE

Products Affected

• ZTALMY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GEFITINIB

Products Affected

• gefitinib

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GILTERITINIB

Products Affected

XOSPATA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GLASDEGIB

Products Affected

• DAURISMO ORAL TABLET 100 MG, 25 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GLATIRAMER

- glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml
- glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GLP1-DULAGLUTIDE

Products Affected

TRULICITY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GLP1-SEMAGLUTIDE

- OZEMPIC
- RYBELSUS

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GLP1-TIRZEPATIDE

Products Affected

MOUNJARO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS. |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GOSERELIN

Products Affected

ZOLADEX

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. |
| Age Restrictions | |
| Prescriber Restrictions | ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. |
| Coverage Duration | STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS. |
| Other Criteria | ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

GUSELKUMAB

- TREMFYA
- TREMFYA PEN SUBCUTANEOUS PEN INJECTOR 200 MG/2 ML

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

Products Affected

• morphine concentrate oral solution

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME. |
| Other Criteria | 1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IBRUTINIB

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ICATIBANT

Products Affected

• icatibant

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING. |
| Age Restrictions | |
| Prescriber Restrictions | HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IDELALISIB

Products Affected

• ZYDELIG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IMATINIB

Products Affected

• imatinib oral tablet 100 mg, 400 mg

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS. |
| Other Criteria | PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IMATINIB SOLUTION

Products Affected

IMKELDI

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS. |
| Other Criteria | PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR. ALL INDICATIONS: UNABLE TO SWALLOW GENERIC IMATINIB TABLETS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IMETELSTAT

Products Affected

• RYTELO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INAVOLISIB

Products Affected

• ITOVEBI ORAL TABLET 3 MG, 9 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INFLIXIMAB

Products Affected

• infliximab

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF |
| | THE FOLLOWING PREFERRED AGENTS: ENBREL, |
| | HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. |
| | PSA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE |
| | FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, |
| | HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, |
| | XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA, |
| | AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC |
| | BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK |
| | INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) TRIAL OF OR |
| | CONTRAINDICATION TO TWO OF THE FOLLOWING |
| | PREFERRED AGENTS: COSENTYX, ENBREL, |
| | HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, |
| | SKYRIZI, TREMFYA, OTEZLA, AND 2) NO CONCURRENT USE |
| | WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |
| | MOLECULES FOR PSO. AS: 1) TRIAL OF OR |
| | CONTRAINDICATION TO TWO OF THE FOLLOWING |
| | PREFERRED AGENTS: COSENTYX, ENBREL, |
| | HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, AND 2) NO |
| | CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR |
| | TARGETED SMALL MOLECULES FOR AS. MODERATE TO |
| | SEVERE CD: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF |
| | THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: |
| | STELARA/SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, |
| | RINVOQ, SKYRIZI, AND 2) NO CONCURRENT USE WITH |
| | ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |
| | MOLECULES FOR CD. UC: 1) TRIAL OF OR CONTRAINDICATION |
| | TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE |
| | AGES ALIGN: STELARA/SELARSDI/YESINTEK, XELJANZ, |
| | HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA, |
| | AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC |
| | BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| | RENEWAL: RA: CONTINUES TO BENEFIT FROM THE |
| | MEDICATION. PSA: 1) CONTINUES TO BENEFIT FROM THE |
| | MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER |
| | |

| PA Criteria | Criteria Details |
|------------------------|--|
| | SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. MODERATE TO SEVERE CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INSULIN SUPPLIES PAYMENT DETERMINATION

- 1ST TIER UNIFINE PENTP 5MM 31G
- 1ST TIER UNIFINE PNTIP 4MM 32G
- 1ST TIER UNIFINE PNTIP 6MM 31G
- 1ST TIER UNIFINE PNTIP 8MM 31G STRL,SINGLE-USE,SHRT
- 1ST TIER UNIFINE PNTP 29GX1/2"
- 1ST TIER UNIFINE PNTP 31GX3/16
- 1ST TIER UNIFINE PNTP 32GX5/32
- ABOUTTIME PEN NEEDLE
- ADVOCATE INS 0.3 ML 30GX5/16"
- ADVOCATE INS 0.3 ML 31GX5/16"
- ADVOCATE INS 0.5 ML 30GX5/16"
- ADVOCATE INS 0.5 ML 31GX5/16"
- ADVOCATE INS 1 ML 31GX5/16"
- ADVOCATE INS SYR 0.3 ML 29GX1/2
- ADVOCATE INS SYR 0.5 ML 29GX1/2
- ADVOCATE INS SYR 1 ML 29GX1/2"
- ADVOCATE INS SYR 1 ML 30GX5/16
- ADVOCATE PEN NDL 12.7MM 29G
- ADVOCATE PEN NEEDLE 32G 4MM
- ADVOCATE PEN NEEDLE 4MM 33G
- ADVOCATE PEN NEEDLES 5MM 31G
- ADVOCATE PEN NEEDLES 8MM 31G
- ALCOHOL 70% SWABS
- ALCOHOL PADS
- ALCOHOL PREP SWABS
- · ALCOHOL WIPES
- AQINJECT PEN NEEDLE 31G 5MM
- AQINJECT PEN NEEDLE 32G 4MM
- ASSURE ID DUO PRO NDL 31G 5MM
- ASSURE ID DUO-SHIELD 30GX3/16"
- ASSURE ID DUO-SHIELD 30GX5/16"
- ASSURE ID INSULIN SAFETY SYRINGE
 1 ML 29 GAUGE X 1/2"
- ASSURE ID PEN NEEDLE 30GX3/16"
- ASSURE ID PEN NEEDLE 30GX5/16"
- ASSURE ID PEN NEEDLE 31GX3/16"
- ASSURE ID PRO PEN NDL 30G 5MM
- ASSURE ID SYR 0.5 ML 29GX1/2" (RX)
- ASSURE ID SYR 0.5 ML 31GX15/64"
- ASSURE ID SYR 1 ML 31GX15/64"

- AUTOSHIELD DUO PEN NDL 30G 5MM
- BD AUTOSHIELD DUO NDL 5MMX30G
- BD ECLIPSE 30GX1/2" SYRINGE
- BD ECLIPSE NEEDLE 30GX1/2" (OTC)
- BD INS SYR 0.3 ML 8MMX31G(1/2)
- BD INS SYR UF 0.3 ML 12.7MMX30G
- BD INS SYR UF 0.5 ML 12.7MMX30G NOT FOR RETAIL SALE
- BD INS SYRN UF 1 ML 12.7MMX30G NOT FOR RETAIL SALE
- BD INS SYRNG UF 0.3 ML 8MMX31G
- BD INS SYRNG UF 0.5 ML 8MMX31G
- BD INSULIN SYR 1 ML 25GX1"
- BD INSULIN SYR 1 ML 25GX5/8"
- BD INSULIN SYR 1 ML 26GX1/2"
- BD INSULIN SYR 1 ML 27GX12.7MM
- BD INSULIN SYR 1 ML 27GX5/8" MICRO-FINE
- BD INSULIN SYRINGE SLIP TIP
- BD INSULIN SYRINGE U-500
- BD LUER-LOK SYRINGE 1 ML
- BD NANO 2 GEN PEN NDL 32G 4MM
- BD SAFETGLD INS 0.3 ML 29G 13MM
- BD SAFETGLD INS 0.5 ML 13MMX29G
- BD SAFETYGLD INS 0.3 ML 31G 8MM
- BD SAFETYGLD INS 0.5 ML 30G 8MM
- BD SAFETYGLD INS 1 ML 29G 13MM
- BD SAFETYGLID INS 1 ML 6MMX31G
- BD SAFETYGLIDE SYRINGE 27GX5/8
- BD SAFTYGLD INS 0.3 ML 6MMX31G
- BD SAFTYGLD INS 0.5 ML 29G 13MM
- BD SAFTYGLD INS 0.5 ML 6MMX31G
- BD SINGLE USE SWAB
- BD UF MICRO PEN NEEDLE 6MMX32G
- BD UF MINI PEN NEEDLE 5MMX31G
- BD UF NANO PEN NEEDLE 4MMX32G
- BD UF ORIG PEN NDL 12.7MMX29G
- BD UF SHORT PEN NEEDLE 8MMX31G
- BD VEO INS 0.3 ML 6MMX31G (1/2)
- BD VEO INS SYRING 1 ML 6MMX31G
- BD VEO INS SYRN 0.3 ML 6MMX31G

- BD VEO INS SYRN 0.5 ML 6MMX31G
- BORDERED GAUZE 2"X2"
- CAREFINE PEN NEEDLE 12.7MM 29G
- CAREFINE PEN NEEDLE 4MM 32G
- CAREFINE PEN NEEDLE 5MM 32G
- CAREFINE PEN NEEDLE 6MM 31G
- CAREFINE PEN NEEDLE 8MM 30G
- CAREFINE PEN NEEDLES 6MM 32G
- CAREFINE PEN NEEDLES 8MM 31G
- CARETOUCH ALCOHOL 70% PREP PAD
- CARETOUCH PEN NEEDLE 29G 12MM
- CARETOUCH PEN NEEDLE 31GX1/4"
- CARETOUCH PEN NEEDLE 31GX3/16"
- CARETOUCH PEN NEEDLE 31GX5/16"
- CARETOUCH PEN NEEDLE 32GX3/16"
- CARETOUCH PEN NEEDLE 32GX5/32"
- CARETOUCH SYR 0.3 ML 31GX5/16"
- CARETOUCH SYR 0.5 ML 30GX5/16"
- CARETOUCH SYR 0.5 ML 31GX5/16"
- CARETOUCH SYR 1 ML 28GX5/16"
- CARETOUCH SYR 1 ML 29GX5/16"
- CARETOUCH SYR 1 ML 30GX5/16"
- CARETOUCH SYR 1 ML 31GX5/16"
- CLICKFINE 31G X 5/16" NEEDLES 8MM, UNIVERSAL
- CLICKFINE PEN NEEDLE 32GX5/32" 32GX4MM, STERILE
- CLICKFINE UNIVERSAL 31G X 1/4" 6MM, STORE BRAND
- COMFORT EZ 0.3 ML 31G 15/64"
- COMFORT EZ 0.5 ML 31G 15/64"
- COMFORT EZ INS 0.3 ML 30GX1/2"
- COMFORT EZ INS 0.3 ML 30GX5/16"
- COMFORT EZ INS 1 ML 31G 15/64"
- COMFORT EZ INS 1 ML 31GX5/16"
- COMFORT EZ INSULIN SYR 0.3 ML
- COMFORT EZ INSULIN SYR 0.5 ML
- COMFORT EZ PEN NEEDLE 12MM 29G
- COMFORT EZ PEN NEEDLES 4MM 32G SINGLE USE, MICRO
- COMFORT EZ PEN NEEDLES 4MM 33G
- COMFORT EZ PEN NEEDLES 5MM 31G MINI
- COMFORT EZ PEN NEEDLES 5MM 32G SINGLE USE,MINI,HRI
- COMFORT EZ PEN NEEDLES 5MM 33G
- COMFORT EZ PEN NEEDLES 6MM 31G

- COMFORT EZ PEN NEEDLES 6MM 32G
- COMFORT EZ PEN NEEDLES 6MM 33G
- COMFORT EZ PEN NEEDLES 8MM 31G SHORT
- COMFORT EZ PEN NEEDLES 8MM 32G
- COMFORT EZ PEN NEEDLES 8MM 33G
- COMFORT EZ PRO PEN NDL 30G 8MM
- COMFORT EZ PRO PEN NDL 31G 4MM
- COMFORT EZ PRO PEN NDL 31G 5MM
- COMFORT EZ SYR 0.3 ML 29GX1/2"
 COMFORT EZ SYR 0.5 ML 28GX1/2"
- COMFORT EZ SYR 0.5 ML 29GX1/2"
- COMFORT EZ SYR 0.5 ML 30GX1/2"
- COMFORT EZ SYR 1 ML 28GX1/2"
- COMFORT EZ SYR 1 ML 29GX1/2"
- COMFORT EZ SYR 1 ML 30GX1/2"
- COMFORT EZ SYR 1 ML 30GX5/16"
- COMFORT POINT PEN NDL 31GX1/3"
- COMFORT POINT PEN NDL 31GX1/6"
- COMFORT TOUCH PEN NDL 31G 4MM
- COMFORT TOUCH PEN NDL 31G 5MM
- COMFORT TOUCH PEN NDL 31G 6MM
- COMFORT TOUCH PEN NDL 31G 8MM
- COMFORT TOUCH PEN NDL 32G 4MM
- COMFORT TOUCH PEN NDL 32G 5MM
- COMFORT TOUCH PEN NDL 32G 6MM
- COMFORT TOUCH PEN NDL 32G 8MM
 COMFORT TOUCH PEN NDL 33G 4MM
- COMFORT TOUCH PEN NDL 33G 6MM
- COMFORT TOUCH PEN NDL 33GX5MM
- CURAD GAUZE PADS 2" X 2"
- CURITY ALCOHOL PREPS 2 PLY,MEDIUM
- CURITY GAUZE SPONGES (12 PLY)-200/BAG
- CURITY GUAZE PADS 1'S(12 PLY)
- DERMACEA 2"X2" GAUZE 12 PLY, USP TYPE VII
- DERMACEA GAUZE 2"X2" SPONGE 8 PLY
- DERMACEA NON-WOVEN 2"X2" SPNGE
- DROPLET 0.3 ML 29G 12.7MM(1/2)
- DROPLET 0.3 ML 30G 12.7MM(1/2)
- DROPLET 0.5 ML 29GX12.5MM(1/2)
- DROPLET 0.5 ML 30GX12.5MM(1/2)
- DROPLET INS 0.3 ML 29GX12.5MM

- DROPLET INS 0.3 ML 30G 8MM(1/2)
- DROPLET INS 0.3 ML 30GX12.5MM
- DROPLET INS 0.3 ML 31G 6MM(1/2)
- DROPLET INS 0.3 ML 31G 8MM(1/2)
- DROPLET INS 0.5 ML 29G 12.7MM
- DROPLET INS 0.5 ML 30G 12.7MM
- DROPLET INS 0.5 ML 30GX6MM(1/2)
- DROPLET INS 0.5 ML 30GX8MM(1/2)
- DROPLET INS 0.5 ML 31GX6MM(1/2)
- DROPLET INS 0.5 ML 31GX8MM(1/2)
- DROPLET INS SYR 0.3 ML 30GX6MM
- DROPLET INS SYR 0.3 ML 30GX8MM
- DROPLET INS SYR 0.3 ML 31GX6MM
- DROPLET INS SYR 0.3 ML 31GX8MM
- DROPLET INS SYR 0.5 ML 30G 8MM
- DROPLET INS SYR 0.5 ML 31G 6MM
- DROPLET INS SYR 0.5 ML 31G 8MM
- DROPLET INS STR 0.3 MIL STO OMINI
- DROPLET INS SYR 1 ML 29GX12.5MM
- DROPLET INS SYR 1 ML 30GX12.5MM
- DROPLET INS SYR 1 ML 30GX6MM
- DROPLET INS SYR 1 ML 30GX8MM
- DROPLET INS SYR 1 ML 31G 6MM
- DROPLET INS SYR 1 ML 31GX6MM
- DROPLET INS SYR 1 ML 31GX8MM
- DROPLET MICRON 34G X 9/64"
- DROPLET PEN NEEDLE 29G 10MM
- DROPLET PEN NEEDLE 29G 12MM
- DROPLET PEN NEEDLE 30G 8MM
- DROPLET PEN NEEDLE 31G 5MM
- DROPLET PEN NEEDLE 31G 6MM
- DROPLET PEN NEEDLE 31G 8MM
- DROPLET PEN NEEDLE 32G 4MM
- DROPLET PEN NEEDLE 32G 5MM
- DROPLET PEN NEEDLE 32G 6MM
- DROPLET PEN NEEDLE 32G 8MM
- DROPSAFE ALCOHOL 70% PREP PADS
- DROPSAFE INS SYR 0.3 ML 31G 6MM
- DROPSAFE INS SYR 0.3 ML 31G 8MM
- DROPSAFE INS SYR 0.5 ML 31G 6MM
- DROPSAFE INS SYR 0.5 ML 31G 8MM
- DROPSAFE INSUL SYR 1 ML 31G 6MM
- DROPSAFE INSUL SYR 1 ML 31G 8MM
- DROPSAFE INSULN 1 ML 29G 12.5MM
- DROPSAFE PEN NEEDLE 31GX1/4"
- DROPSAFE PEN NEEDLE 31GX3/16"
- DROPSAFE PEN NEEDLE 31GX5/16"
- DRUG MART ULTRA COMFORT SYR

- EASY CMFT SFTY PEN NDL 31G 5MM
- EASY CMFT SFTY PEN NDL 31G 6MM
- EASY CMFT SFTY PEN NDL 32G 4MM
- EASY COMFORT 0.3 ML 31G 1/2"
- EASY COMFORT 0.3 ML 31G 5/16"
- EASY COMFORT 0.3 ML SYRINGE
- EASY COMFORT 0.5 ML 30GX1/2"
- EASY COMFORT 0.5 ML 31GX5/16"
- EASY COMFORT 0.5 ML 32GX5/16"
- EASY COMFORT 0.5 ML SYRINGE
- EASY COMFORT 1 ML 31GX5/16"
- EASY COMFORT 1 ML 32GX5/16"
- EASY COMFORT ALCOHOL 70% PAD
- EASY COMFORT INSULIN 1 ML SYR
- EASY COMFORT PEN NDL 31GX1/4"
- EASY COMFORT PEN NDL 31GX3/16"
- EASY COMFORT PEN NDL 31GX5/16"
- EASY COMFORT PEN NDL 32GX5/32"
- EASY COMFORT PEN NDL 33G 4MM
- EASY COMFORT PEN NDL 33G 5MM
- EASY COMFORT PEN NDL 33G 6MM
- EASY COMFORT SYR 0.5 ML 29G 8MM
- EASY COMFORT SYR 1 ML 29G 8MM
- EASY COMFORT SYR 1 ML 30GX1/2"
- EASY GLIDE INS 0.3 ML 31GX6MM
- EASY GLIDE INS 0.5 ML 31GX6MM
 EASY GLIDE INS 1 ML 31GX6MM
- EASY GLIDE PEN NEEDLE 4MM 33G
- EASY TOUCH 0.3 ML SYR 30GX1/2"
- EASY TOUCH 0.5 ML SYR 27GX1/2"
- EASY TOUCH 0.5 ML SYR 29GX1/2"
- EASY TOUCH 0.5 ML SYR 30GX1/2"
- EASY TOUCH 0.5 ML SYR 30GX5/16
- EASY TOUCH 1 ML SYR 27GX1/2"
- EASY TOUCH 1 ML SYR 29GX1/2"
 EASY TOUCH 1 ML SYR 30GX1/2"
- EASY TOUCH ALCOHOL 70% PADS GAMMA-STERILIZED
- EASY TOUCH FLIPLOK 1 ML 27GX0.5
- EASY TOUCH INSULIN 1 ML 29GX1/2
- EASY TOUCH INSULIN 1 ML 30GX1/2
- EASY TOUCH INSULIN SYR 0.3 ML
- EASY TOUCH INSULIN SYR 0.5 ML
- EASY TOUCH INSULIN SYR 1 ML
- EASY TOUCH INSULIN SYR 1 ML RETRACTABLE
- EASY TOUCH INSULN 1 ML 29GX1/2"

- EASY TOUCH INSULN 1 ML 30GX1/2"
- EASY TOUCH INSULN 1 ML 30GX5/16
- EASY TOUCH INSULN 1 ML 31GX5/16
- EASY TOUCH LUER LOK INSUL 1 ML
- EASY TOUCH PEN NEEDLE 29GX1/2"
- EASY TOUCH PEN NEEDLE 30GX5/16
- EASY TOUCH PEN NEEDLE 31GX1/4"
- EASY TOUCH PEN NEEDLE 31GX3/16
- EASY TOUCH PEN NEEDLE 31GX5/16
- EASY TOUCH PEN NEEDLE 32GX1/4"
- EASY TOUCH PEN NEEDLE 32GX3/16
- EASY TOUCH PEN NEEDLE 32GX5/32
- EASY TOUCH SAF PEN NDL 29G 5MM
- EASY TOUCH SAF PEN NDL 29G 8MM
- EASY TOUCH SAF PEN NDL 30G 5MM
- EASY TOUCH SAF PEN NDL 30G 8MM
- EASY TOUCH SYR 0.5 ML 28G 12.7MM
- EASY TOUCH SYR 0.5 ML 29G 12.7MM
- EASY TOUCH SYR 1 ML 27G 16MM
- EASY TOUCH SYR 1 ML 28G 12.7MM
- EASY TOUCH SYR 1 ML 29G 12.7MM
- EASY TOUCH UNI-SLIP SYR 1 ML
- EASYTOUCH SAF PEN NDL 30G 6MM
- EMBRACE PEN NEEDLE 29G 12MM
- EMBRACE PEN NEEDLE 30G 5MM
- EMBRACE PEN NEEDLE 30G 8MM
- EMBRACE PEN NEEDLE 31G 5MM
- EMBRACE PEN NEEDLE 31G 6MM
- EMBRACE PEN NEEDLE 31G 8MM
- EMBRACE PEN NEEDLE 32G 4MM
- EQL INSULIN 0.3 ML SYRINGE SHORT **NEEDLE**
- EQL INSULIN 0.5 ML SYRINGE SHORT **NEEDLE**
- EQL INSULIN 1 ML SYRINGE SHORT **NEEDLE**
- FIFTY50 INS SYR 1 ML 31GX5/16" SHORT NEEDLE (OTC)
- FIFTY50 PEN 31G X 3/16" NEEDLE (OTC)
- FP INSULIN 1 ML SYRINGE
- FREESTYLE PREC 0.5 ML 30GX5/16
- FREESTYLE PREC 0.5 ML 31GX5/16
- FREESTYLE PREC 1 ML 30GX5/16"
- FREESTYLE PREC 1 ML 31GX5/16"
- GAUZE PAD TOPICAL BANDAGE 2 X 2

- GNP ULT C 0.3 ML 29GX1/2" (1/2) 1/2
- GNP ULTRA COMFORT 0.5 ML SYR
- GNP ULTRA COMFORT 1 ML SYRINGE
- GNP ULTRA COMFORT 3/10 ML SYR
- HEALTHWISE INS 0.3 ML 30GX5/16"
- HEALTHWISE INS 0.3 ML 31GX5/16"
- HEALTHWISE INS 0.5 ML 30GX5/16"
- HEALTHWISE INS 0.5 ML 31GX5/16"
- HEALTHWISE INS 1 ML 30GX5/16"
- HEALTHWISE INS 1 ML 31GX5/16"
- HEALTHWISE PEN NEEDLE 31G 5MM
- HEALTHWISE PEN NEEDLE 31G 8MM
- HEALTHWISE PEN NEEDLE 32G 4MM
- **HEALTHY ACCENTS PENTIP 4MM 32G**
- **HEALTHY ACCENTS PENTIP 5MM 31G**
- **HEALTHY ACCENTS PENTIP 6MM 31G**
- **HEALTHY ACCENTS PENTIP 8MM 31G**
- **HEALTHY ACCENTS PENTP 12MM 29G**
- HEB INCONTROL ALCOHOL 70% PADS
- **INCONTROL PEN NEEDLE 12MM 29G**
- **INCONTROL PEN NEEDLE 4MM 32G**
- **INCONTROL PEN NEEDLE 5MM 31G**
- **INCONTROL PEN NEEDLE 6MM 31G**
- **INCONTROL PEN NEEDLE 8MM 31G**
- INSULIN SYR 0.3 ML 31GX1/4(1/2)
- INSULIN SYRIN 0.5 ML 28GX1/2" (OTC)
- INSULIN SYRIN 0.5 ML 29GX1/2" (OTC)
- INSULIN SYRIN 0.5 ML 30GX1/2" (RX)
- INSULIN SYRIN 0.5 ML 30GX5/16" SHORT NEEDLE (OTC)
- INSULIN SYRING 0.5 ML 27G 1/2" OUTER
- **INSULIN SYRINGE 0.3 ML**
- INSULIN SYRINGE 0.3 ML 31GX1/4
- **INSULIN SYRINGE 0.5 ML**
- INSULIN SYRINGE 0.5 ML 31GX1/4
- **INSULIN SYRINGE 1 ML**
- INSULIN SYRINGE 1 ML 27G 1/2" **INNER**
- **INSULIN SYRINGE 1 ML 27G 16MM**
- INSULIN SYRINGE 1 ML 28GX1/2" (OTC)
- INSULIN SYRINGE 1 ML 30GX1/2" (RX)
- INSULIN SYRINGE 1 ML 30GX5/16" SHORT NEEDLE (OTC)
- INSULIN SYRINGE 1 ML 31GX1/4"

- INSULIN SYRINGE-NEEDLE U-100 SYRINGE 0.3 ML 29 GAUGE, 1 ML 29 GAUGE X 1/2", 1/2 ML 28 GAUGE
- INSUPEN 30G ULTRAFIN NEEDLE
- INSUPEN 31G ULTRAFIN NEEDLE
- INSUPEN 32G 6MM PEN NEEDLE
- INSUPEN 32G 8MM PEN NEEDLE
- INSUPEN PEN NEEDLE 29GX12MM
- INSUPEN PEN NEEDLE 31GX3/16"
- INSUPEN PEN NEEDLE 32GX4MM
- INSUPEN PEN NEEDLE 33GX4MM
- IV ANTISEPTIC WIPES
- KENDALL ALCOHOL 70% PREP PAD
- LISCO SPONGES 100/BAG
- LITE TOUCH 31GX1/4" PEN NEEDLE
- LITE TOUCH INSULIN 0.5 ML SYR
- LITE TOUCH INSULIN 1 ML SYR
- LITE TOUCH INSULIN SYR 1 ML
- LITE TOUCH PEN NEEDLE 29G
- LITE TOUCH PEN NEEDLE 31G
- LITETOUCH INS 0.3 ML 29GX1/2"
- LITETOUCH INS 0.3 ML 30GX5/16"
- LITETOUCH INS 0.3 ML 31GX5/16"
- LITETOUCH INS 0.5 ML 31GX5/16"
- LITETOUCH SYR 0.5 ML 28GX1/2"
- LITETOUCH SYR 0.5 ML 29GX1/2"
- LITETOUCH SYR 0.5 ML 30GX5/16"
- LITETOUCH SYRIN 1 ML 28GX1/2"
- LITETOUCH SYRIN 1 ML 29GX1/2"
- LITETOUCH SYRIN 1 ML 30GX5/16"
- MAGELLAN INSUL SYRINGE 0.3 ML
- MAGELLAN INSUL SYRINGE 0.5 ML
- MAGELLAN INSULIN SYR 0.3 ML
- MAGELLAN INSULIN SYR 0.5 ML
- MAGELLAN INSULIN SYRINGE 1 ML
- MAXI-COMFORT INS 0.5 ML 28G
- MAXI-COMFORT INS 1 ML 28GX1/2"
- MAXICOMFORT II PEN NDL 31GX6MM
- MAXICOMFORT INS 0.5 ML 27GX1/2"
- MAXICOMFORT INS 1 ML 27GX1/2"
- MAXICOMFORT PEN NDL 29G X 5MM
- MAXICOMFORT PEN NDL 29G X 8MM
- MICRODOT PEN NEEDLE 31GX6MM
- MICRODOT PEN NEEDLE 32GX4MM
- MICRODOT PEN NEEDLE 33GX4MM
- MICRODOT READYGARD NDL 31G 5MM OUTER

- MINI PEN NEEDLE 32G 4MM
- MINI PEN NEEDLE 32G 5MM
- MINI PEN NEEDLE 32G 6MM
- MINI PEN NEEDLE 32G 8MM
- MINI PEN NEEDLE 33G 4MM
- MINI PEN NEEDLE 33G 5MM
- MINI PEN NEEDLE 33G 6MM
- MINI ULTRA-THIN II PEN NDL 31G STERILE
- MONOJECT 0.5 ML SYRN 28GX1/2"
- MONOJECT 1 ML SYRN 27X1/2"
- MONOJECT 1 ML SYRN 28GX1/2" (OTC)
- MONOJECT INSUL SYR U100 (OTC)
- MONOJECT INSUL SYR U100 .5ML,29GX1/2" (OTC)
- MONOJECT INSUL SYR U100 0.5 ML CONVERTS TO 29G (OTC)
- MONOJECT INSUL SYR U100 1 ML
- MONOJECT INSUL SYR U100 1 ML 3'S, 29GX1/2" (OTC)
- MONOJECT INSUL SYR U100 1 ML W/O NEEDLE (OTC)
- MONOJECT INSULIN SYR 0.3 ML
- MONOJECT INSULIN SYR 0.3 ML (OTC)
- MONOJECT INSULIN SYR 0.5 ML
- MONOJECT INSULIN SYR 0.5 ML (OTC)
- MONOJECT INSULIN SYR 1 ML 3'S (OTC)
- MONOJECT INSULIN SYR U-100
- MONOJECT SYRINGE 0.3 ML
- MONOJECT SYRINGE 0.5 ML
- MONOJECT SYRINGE 1 ML
- NANO 2 GEN PEN NEEDLE 32G 4MM
- NOVOFINE 30
- NOVOFINE 32G NEEDLES
- NOVOFINE PLUS PEN NDL 32GX1/6"
- NOVOTWIST NEEDLE 32G 5MM
- PC UNIFINE PENTIPS 8MM NEEDLE SHORT
- PEN NEEDLE 30G 5MM OUTER
- PEN NEEDLE 30G 8MM INNER
- PEN NEEDLE 30G X 5/16"
- PEN NEEDLE, DIABETIC NEEDLE 29 GAUGE X 1/2"
- PEN NEEDLES 12MM 29G 29GX12MM,STRL
- PEN NEEDLES 4MM 32G

- PEN NEEDLES 6MM 31G 31GX6MM, STRI.
- PEN NEEDLES 8MM 31G 31GX8MM,STRL,SHORT (OTC)
- PENTIPS PEN NEEDLE 29G 1/2"
- PENTIPS PEN NEEDLE 31G 1/4"
- PENTIPS PEN NEEDLE 31GX3/16" MINI, 5MM
- PENTIPS PEN NEEDLE 31GX5/16" SHORT, 8MM
- PENTIPS PEN NEEDLE 32G 1/4"
- PENTIPS PEN NEEDLE 32GX5/32" 4MM
- PIP PEN NEEDLE 31G X 5MM
- PIP PEN NEEDLE 32G X 4MM
- PREVENT PEN NEEDLE 31GX1/4"
- PREVENT PEN NEEDLE 31GX5/16"
- PRO COMFORT 0.5 ML 30GX1/2"
- PRO COMFORT 0.5 ML 30GX5/16"
- PRO COMFORT 0.5 ML 31GX5/16"
- PRO COMFORT 1 ML 30GX1/2"
- PRO COMFORT 1 ML 30GX5/16"
- PRO COMFORT 1 ML 31GX5/16"
- PRO COMFORT ALCOHOL 70% PADS
- PRO COMFORT PEN NDL 31GX5/16"
- PRO COMFORT PEN NDL 32G X 1/4"
- PRO COMFORT PEN NDL 4MM 32G
- PRO COMFORT PEN NDL 5MM 32G
- PRODIGY INS SYR 1 ML 28GX1/2"
- PRODIGY SYRNG 0.5 ML 31GX5/16"
- PRODIGY SYRNGE 0.3 ML 31GX5/16"
- PURE CMFT SFTY PEN NDL 31G 5MM
- PURE CMFT SFTY PEN NDL 31G 6MM
- PURE CMFT SFTY PEN NDL 32G 4MM
- PURE COMFORT ALCOHOL 70% PADS
- PURE COMFORT PEN NDL 32G 4MM
- PURE COMFORT PEN NDL 32G 5MM
- PURE COMFORT PEN NDL 32G 6MM
- PURE COMFORT PEN NDL 32G 8MM
- RAYA SURE PEN NEEDLE 29G 12MM
- RAYA SURE PEN NEEDLE 31G 4MM
- RAYA SURE PEN NEEDLE 31G 5MM
- RAYA SURE PEN NEEDLE 31G 6MM
- RELI-ON INSULIN 0.5 ML SYR
- RELI-ON INSULIN 1 ML SYR
- RELION INS SYR 0.3 ML 31GX6MM
- RELION INS SYR 0.5 ML 31GX6MM
- RELION INS SYR 1 ML 31GX15/64"

- RELION MINI PEN 31G X 1/4" NDL
- SAFESNAP INS SYR UNITS-100 0.3 ML 30GX5/16",10X10
- SAFESNAP INS SYR UNITS-100 0.5 ML 29GX1/2",10X10
- SAFESNAP INS SYR UNITS-100 0.5 ML 30GX5/16",10X10
- SAFESNAP INS SYR UNITS-100 1 ML 28GX1/2",10X10
- SAFESNAP INS SYR UNITS-100 1 ML 29GX1/2",10X10
- SAFETY PEN NEEDLE 31G 4MM
- SAFETY PEN NEEDLE 5MM X 31G
- SAFETY SYRINGE 0.5 ML 30G 1/2"
- SECURESAFE PEN NDL 30GX5/16" OUTER
- SECURESAFE SYR 0.5 ML 29G 1/2" OUTER
- SECURESAFE SYRNG 1 ML 29G 1/2" OUTER
- SKY SAFETY PEN NEEDLE 30G 5MM
- SKY SAFETY PEN NEEDLE 30G 8MM
- SM ULT CFT 0.3 ML 31GX5/16(1/2)
- STERILE PADS 2" X 2"
- SURE CMFT SFTY PEN NDL 31G 6MM
- SURE CMFT SFTY PEN NDL 32G 4MM
- SURE COMFORT 0.5 ML SYRINGE
- SURE COMFORT 1 ML SYRINGE
- SURE COMFORT 3/10 ML SYRINGE
- SURE COMFORT 3/10 ML SYRINGE INSULIN SYRINGE
- SURE COMFORT 30G PEN NEEDLE
- SURE COMFORT ALCOHOL PREP PADS
- SURE COMFORT INS 0.3 ML 31GX1/4
- SURE COMFORT INS 0.5 ML 31GX1/4
- SURE COMFORT INS 1 ML 31GX1/4"
- SURE COMFORT PEN NDL 29GX1/2"
 12.7MM
- SURE COMFORT PEN NDL 31G 5MM
- SURE COMFORT PEN NDL 31G 8MM
- SURE COMFORT PEN NDL 32G 4MM
- SURE COMFORT PEN NDL 32G 6MM
- SURE-FINE PEN NEEDLES 12.7MM
- SURE-FINE PEN NEEDLES 5MM
- SURE-FINE PEN NEEDLES 8MM
- SURE-JECT INSU SYR U100 0.3 ML
- SURE-JECT INSU SYR U100 0.5 ML

- SURE-JECT INSU SYR U100 1 ML
- SURE-JECT INSUL SYR U100 1 ML
- SURE-JECT INSULIN SYRINGE 1 ML
- SURE-PREP ALCOHOL PREP PADS
- TECHLITE 0.3 ML 29GX12MM (1/2)
- TECHLITE 0.3 ML 30GX8MM (1/2)
- TECHLITE 0.3 ML 31GX6MM (1/2)
- TECHLITE 0.3 ML 31GX8MM (1/2)
- TECHLITE 0.5 ML 30GX12MM (1/2)
- TECHLITE 0.5 ML 30GX8MM (1/2)
- TECHLITE 0.5 ML 31GX6MM (1/2)
- TECHLITE 0.5 ML 31GX8MM (1/2)
- TECHLITE INS SYR 1 ML 29GX12MM
- TECHLITE INS SYR 1 ML 30GX12MM
- TECHLITE INS SYR 1 ML 30GX8MM
- TECHLITE INS SYR 1 ML 31GX6MM
- TECHLITE INS SYR 1 ML 31GX8MM
- TECHLITE INS STR 1 ML STGX8MM
 TECHLITE PEN NEEDLE 29GX1/2"
- TECHLITE PEN NEEDLE 29GX3/8"
- TECHLITE PEN NEEDLE 31GX1/4"
- TECHLITE PEN NEEDLE 31GX3/16"
- TECHLITE PEN NEEDLE 31GX5/16"
- TECHLITE DENINGED E 22CV1/4
- TECHLITE PEN NEEDLE 32GX1/4"
- TECHLITE PEN NEEDLE 32GX5/16"
- TECHLITE PEN NEEDLE 32GX5/32"
- TECHLITE PLUS PEN NDL 32G 4MM
- TERUMO INS SYRINGE U100-1 ML
- TERUMO INS SYRINGE U100-1/2 ML
- TERUMO INS SYRINGE U100-1/3 ML
- TERUMO INS SYRNG U100-1/2 ML
- THINPRO INS SYRIN U100-0.3 ML
- THINPRO INS SYRIN U100-0.5 ML
- THINPRO INS SYRIN U100-1 ML
- TOPCARE CLICKFINE 31G X 1/4"
- TOPCARE CLICKFINE 31G X 5/16"
- TOPCARE ULTRA COMFORT SYRINGE
- TRUE CMFRT PRO 0.5 ML 30G 5/16"
- TRUE CMFRT PRO 0.5 ML 31G 5/16"
- TRUE CMFRT PRO 0.5 ML 32G 5/16"
- TRUE CMFT SFTY PEN NDL 31G 5MM
- TRUE CMFT SFTY PEN NDL 31G 6MM
- TRUE CMFT SFTY PEN NDL 32G 4MM
- TRUE COMFORT 0.5 ML 30G 1/2"
- TRUE COMFORT 0.5 ML 30G 5/16"
- TRUE COMFORT 0.5 ML 31G 5/16"
- TRUE COMFORT 0.5 ML 31GX5/16"
- TRUE COMFORT 1 ML 31GX5/16"

- TRUE COMFORT ALCOHOL 70% PADS
- TRUE COMFORT PEN NDL 31G 8MM
- TRUE COMFORT PEN NDL 31GX5MM
- TRUE COMFORT PEN NDL 31GX6MM
- TRUE COMFORT PEN NDL 32G 5MM
- TRUE COMFORT PEN NDL 32G 6MM
- TRUE COMFORT PEN NDL 32GX4MM
- TRUE COMFORT PEN NDL 33G 4MM
- TRUE COMFORT PEN NDL 33G 5MM
- TRUE COMFORT PEN NDL 33G 6MM
- TRUE COMFORT PRO 1 ML 30G 1/2"
- TRUE COMFORT PRO 1 ML 30G 5/16"
- TRUE COMPORT I RO 1 ML 300 3/10
- TRUE COMFORT PRO 1 ML 31G 5/16"
- TRUE COMFORT PRO 1 ML 32G 5/16"
- TRUE COMFORT PRO ALCOHOL PADS
- TRUE COMFORT SFTY 1 ML 30G 1/2"
- TRUE COMFRT PRO 0.5 ML 30G 1/2"
- TRUE COMFRT SFTY 1 ML 30G 5/16"
- TRUE COMFRT SFTY 1 ML 31G 5/16"
- TRUE COMFRT SFTY 1 ML 32G 5/16"
- TRUEPLUS PEN NEEDLE 29GX1/2"
- TRUEPLUS PEN NEEDLE 31G X 1/4"
- TRUEPLUS PEN NEEDLE 31GX3/16"
- TRUEPLUS PEN NEEDLE 31GX5/16"
- TRUEPLUS PEN NEEDLE 32GX5/32"
- TRUEPLUS SYR 0.3 ML 29GX1/2"
- TRUEPLUS SYR 0.3 ML 30GX5/16"
- TRUEPLUS SYR 0.3 ML 31GX5/16"
- TRUEPLUS SYR 0.5 ML 28GX1/2"
- TRUEPLUS SYR 0.5 ML 29GX1/2"
- TRUEPLUS SYR 0.5 ML 30GX5/16"
- TRUEPLUS SYR 0.5 ML 31GX5/16"
- TRUEPLUS SYR 1 ML 28GX1/2"
- TRUEPLUS SYR 1 ML 29GX1/2"
- TRUEPLUS SYR 1 ML 30GX5/16"
- TRUEPLUS SYR 1 ML 31GX5/16"
- ULTICAR INS 0.3 ML 31GX1/4(1/2)
- ULTICARE INS 1 ML 31GX1/4"
- ULTICARE INS SYR 0.3 ML 30G 8MM
- ULTICARE INS SYR 0.3 ML 31G 6MM
- ULTICARE INS SYR 0.3 ML 31G 8MM
- ULTICARE INS SYR 0.5 ML 31G 6MM
- ULTICARE INS SYR 0.5 ML 31G 8MM (OTC)
- ULTICARE INS SYR 1 ML 30GX1/2"
- ULTICARE PEN NEEDLE 31GX3/16"
- ULTICARE PEN NEEDLE 6MM 31G

- ULTICARE PEN NEEDLE 8MM 31G
- ULTICARE PEN NEEDLES 12MM 29G
- ULTICARE PEN NEEDLES 4MM 32G MICRO, 32GX4MM
- ULTICARE PEN NEEDLES 6MM 32G
- ULTICARE SAFE PEN NDL 30G 8MM
- ULTICARE SAFE PEN NDL 5MM 30G
- ULTICARE SYR 0.3 ML 29G 12.7MM
- ULTICARE SYR 0.3 ML 30GX1/2"
- ULTICARE SYR 0.3 ML 31GX5/16" SHORT NDL
- ULTICARE SYR 0.5 ML 30GX1/2"
- ULTICARE SYR 0.5 ML 31GX5/16" SHORT NDL
- ULTICARE SYR 1 ML 31GX5/16"
- ULTIGUARD SAFE 1 ML 30G 12.7MM
- ULTIGUARD SAFE0.3 ML 30G 12.7MM
- ULTIGUARD SAFE0.5 ML 30G 12.7MM
- ULTIGUARD SAFEPACK 1 ML 31G 8MM
- ULTIGUARD SAFEPACK 29G 12.7MM
- ULTIGUARD SAFEPACK 31G 5MM
- ULTIGUARD SAFEPACK 31G 6MM
- ULTIGUARD SAFEPACK 31G 8MM
- ULTIGUARD SAFEPACK 32G 4MM
- ULTIGUARD SAFEPACK 32G 6MM
- ULTIGUARD SAFEPK 0.3 ML 31G 8MM
- ULTIGUARD SAFEPK 0.5 ML 31G 8MM
- ULTILET ALCOHOL STERL SWAB
- ULTILET INSULIN SYRINGE 0.3 ML
- ULTILET INSULIN SYRINGE 0.5 ML
- ULTILET INSULIN SYRINGE 1 ML
- ULTILET PEN NEEDLE
- ULTILET PEN NEEDLE 4MM 32G
- ULTRA COMFORT 0.3 ML SYRINGE
- ULTRA COMFORT 0.5 ML 28GX1/2" CONVERTS TO 29G
- ULTRA COMFORT 0.5 ML 29GX1/2"
- ULTRA COMFORT 0.5 ML SYRINGE
- ULTRA COMFORT 1 ML 31GX5/16"
- ULTRA COMFORT 1 ML SYRINGE
- ULTRA FLO 0.3 ML 30G 1/2" (1/2)
- ULTRA FLO 0.3 ML 30G 5/16"(1/2)
- ULTRA FLO 0.3 ML 31G 5/16"(1/2)
- ULTRA FLO PEN NEEDLE 31G 5MM
- ULTRA FLO PEN NEEDLE 31G 8MM
- ULTRA FLO PEN NEEDLE 32G 4MM
- ULTRA FLO PEN NEEDLE 33G 4MM

- ULTRA FLO PEN NEEDLES 12MM 29G
- ULTRA FLO SYR 0.3 ML 29GX1/2"
- ULTRA FLO SYR 0.3 ML 30G 5/16"
- ULTRA FLO SYR 0.3 ML 31G 5/16"
- ULTRA FLO SYR 0.5 ML 29G 1/2"
- ULTRA THIN PEN NDL 32G X 4MM
- ULTRA-FINE 0.3 ML 30G 12.7MM
- ULTRA-FINE 0.3 ML 31G 6MM (1/2)
- ULTRA-FINE 0.3 ML 31G 8MM (1/2)
- ULTRA-FINE 0.5 ML 30G 12.7MM
- ULTRA-FINE INS SYR 1 ML 31G 8MM
- ULTRA-FINE PEN NDL 29G 12.7MM
- ULTRA-FINE PEN NEEDLE 32G 6MM
- ULTRA-FINE SYR 0.5 ML 31G 8MM
- ULTRA-FINE SYR 1 ML 30G 12.7MM
 ULTRA-THIN II 1 ML 31GX5/16"
- ULTRA-THIN II INS 0.3 ML 30G
- ULTRA-THIN II INS 0.3 ML 31G
- ULTRA-THIN II INS 0.5 ML 29G
- ULTRA-THIN II INS 0.5 ML 30G
- ULTRA-THIN II INS 0.5 ML 31G
- ULTRA-THIN II INS SYR 1 ML 29G
- ULTRA-THIN II INS SYR 1 ML 30G
- ULTRA-THIN II PEN NDL 29GX1/2"
- ULTRA-THIN II PEN NDL 31GX5/16
- ULTRACARE INS 0.3 ML 30GX5/16"
- ULTRACARE INS 0.3 ML 31GX5/16"
- ULTRACARE INS 0.5 ML 30GX1/2"
- ULTRACARE INS 0.5 ML 30GX5/16"
- ULTRACARE INS 0.5 ML 31GX5/16"
- ULTRACARE INS 1 ML 30G X 5/16"
- ULTRACARE INS 1 ML 30GX1/2"
- ULTRACARE INS 1 ML 31G X 5/16"
- ULTRACARE PEN NEEDLE 31GX1/4"
- ULTRACARE PEN NEEDLE 31GX3/16"
- ULTRACARE PEN NEEDLE 31GX5/16"
- ULTRACARE PEN NEEDLE 32GX1/4"
- ULTRACARE PEN NEEDLE 32GX3/16"
- ULTRACARE PEN NEEDLE 32GX5/32"
- ULTRACARE PEN NEEDLE 33GX5/32"
- UNIFINE OTC PEN NEEDLE 31G 5MM
- UNIFINE OTC PEN NEEDLE 32G 4MM
- UNIFINE PEN NEEDLE 32G 4MM
- UNIFINE PENTIPS 12MM 29G 29GX12MM, STRL
- UNIFINE PENTIPS 31GX3/16" 31GX5MM,STRL,MINI

- UNIFINE PENTIPS 32GX1/4"
- UNIFINE PENTIPS 32GX5/32" 32GX4MM, STRL, NANO
- UNIFINE PENTIPS 33GX5/32"
- UNIFINE PENTIPS 6MM 31G
- UNIFINE PENTIPS MAX 30GX3/16"
- UNIFINE PENTIPS NEEDLES 29G
- UNIFINE PENTIPS PLUS 29GX1/2" 12MM
- UNIFINE PENTIPS PLUS 30GX3/16"
- UNIFINE PENTIPS PLUS 31GX1/4" ULTRA SHORT, 6MM
- UNIFINE PENTIPS PLUS 31GX3/16" MINI
- UNIFINE PENTIPS PLUS 31GX5/16" SHORT
- UNIFINE PENTIPS PLUS 32GX5/32"
- UNIFINE PENTIPS PLUS 33GX5/32"
- UNIFINE PROTECT 30G 5MM
- UNIFINE PROTECT 30G 8MM
- UNIFINE PROTECT 32G 4MM
- UNIFINE SAFECONTROL 30G 5MM
- UNIFINE SAFECONTROL 30G 8MM
- UNIFINE SAFECONTROL 31G 5MM
- UNIFINE SAFECONTROL 31G 6MM
- UNIFINE SAFECONTROL 31G 8MM
- UNIFINE SAFECONTROL 32G 4MM
- UNIFINE ULTRA PEN NDL 31G 5MM

- UNIFINE ULTRA PEN NDL 31G 6MM
- UNIFINE ULTRA PEN NDL 31G 8MM
- UNIFINE ULTRA PEN NDL 32G 4MM
- VANISHPOINT 0.5 ML 30GX1/2" SY OUTER
- VANISHPOINT INS 1 ML 30GX3/16"
- VANISHPOINT U-100 29X1/2 SYR
- VERIFINE INS SYR 1 ML 29G 1/2"
- VERIFINE PEN NEEDLE 29G 12MM
- VERIFINE PEN NEEDLE 31G 5MM
- VERIFINE PEN NEEDLE 31G X 6MM
- VERIFINE PEN NEEDLE 31G X 8MM
- VERIFINE PEN NEEDLE 32G 6MM
- VERIFINE PEN NEEDLE 32G X 4MM
- VERIFINE PEN NEEDLE 32G X 5MM
- VERIFINE PLUS PEN NDL 31G 5MM
- VERIFINE PLUS PEN NDL 31G 8MM
- VERIFINE PLUS PEN NDL 32G 4MM
- VERIFINE PLUS PEN NDL 32G 4MM-SHARPS CONTAINER
- VERIFINE SYRING 0.5 ML 29G 1/2"
- VERIFINE SYRING 1 ML 31G 5/16"
- VERIFINE SYRNG 0.3 ML 31G 5/16"
- VERIFINE SYRNG 0.5 ML 31G 5/16"
- VERSALON ALL PURPOSE SPONGE 25'S,N-STERILE,3PLY
- WEBCOL ALCOHOL PREPS 20'S,LARGE

| PA Criteria | Criteria Details |
|------------------------------------|------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | LIFETIME |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | ONLY COVERED UNDER PART D WHEN USED CONCURRENTLY WITH INSULIN. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INTERFERON FOR MS-AVONEX

Products Affected

KIT

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE
- AVONEX PEN 30 MCG/0.5 ML

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INTERFERON FOR MS-BETASERON

Products Affected

• BETASERON SUBCUTANEOUS KIT

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INTERFERON FOR MS-PLEGRIDY

Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE

125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INTERFERON GAMMA-1B

Products Affected

ACTIMMUNE

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR HEMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IPILIMUMAB

Products Affected

YERVOY

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO |
| Other Criteria | RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IVACAFTOR

Products Affected

KALYDECO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS |
| Age Restrictions | |
| Prescriber Restrictions | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT |
| Coverage Duration | INITIAL: 12 MONTHS. RENEWAL: LIFETIME |
| Other Criteria | CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: IMPROVEMENT IN CLINICAL STATUS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IVOSIDENIB

Products Affected

TIBSOVO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

IXAZOMIB

Products Affected

NINLARO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LANREOTIDE

- lanreotide subcutaneous syringe 120 mg/0.5 ml
- SOMATULINE DEPOT SUBCUTANEOUS SYRINGE 60 MG/0.2 ML, 90 MG/0.3 ML

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP-NETS, CARCINOID SYNDROME: 12 MOS. |
| Other Criteria | ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LAPATINIB

Products Affected

• lapatinib

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LAROTRECTINIB

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | VITRAKVI ORAL SOLUTION: 1) TRIAL OF VITRAKVI CAPSULES, OR 2) UNABLE TO TAKE CAPSULE FORMULATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LAZERTINIB

Products Affected

• LAZCLUZE ORAL TABLET 240 MG, 80 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LEDIPASVIR-SOFOSBUVIR

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. |
| Other Criteria | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANAVIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LENALIDOMIDE

Products Affected

• lenalidomide

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LENVATINIB

Products Affected

LENVIMA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LETERMOVIR

Products Affected

• PREVYMIS ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS. |
| Other Criteria | HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LEUPROLIDE

Products Affected

• leuprolide subcutaneous kit

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | PROSTATE CANCER: 12 MONTHS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LEUPROLIDE DEPOT

Products Affected

• leuprolide (3 month)

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LEUPROLIDE-ELIGARD

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LEUPROLIDE-LUPRON DEPOT

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. |
| Coverage Duration | PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS. |
| Other Criteria | INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

LEUPROLIDE-LUPRON DEPOT-PED

- LUPRON DEPOT-PED (3 MONTH)
- LUPRON DEPOT-PED INTRAMUSCULAR SYRINGE KIT

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. |
| Age Restrictions | |
| Prescriber Restrictions | CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

L-GLUTAMINE

Products Affected

• glutamine (sickle cell)

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST |
| Coverage Duration | INITIAL: 12 MONTHS. RENEWAL: LIFETIME. |
| Other Criteria | SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LIDOCAINE OINTMENT

Products Affected

• lidocaine topical ointment

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LIDOCAINE PATCH

- ZTLIDO
- dermacinrx lidocan 5% patch outerlidocaine topical adhesive patch,medicated 5
- lidocan iii

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | 1) PAIN ASSOCIATED WITH POST-HERPETIC NEURALGIA, 2) NEUROPATHY DUE TO DIABETES MELLITUS, 3) CHRONIC BACK PAIN, OR 4) OSTEOARTHRITIS OF THE KNEE OR HIP. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LIDOCAINE PRILOCAINE

Products Affected

• lidocaine-prilocaine topical cream

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LONCASTUXIMAB TESIRINE-LPYL

Products Affected

ZYNLONTA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LORLATINIB

Products Affected

• LORBRENA ORAL TABLET 100 MG, 25 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LOTILANER

Products Affected

• XDEMVY

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | DEMODEX BLEPHARITIS: 18 YEARS OF AGE OR OLDER |
| Prescriber Restrictions | |
| Coverage Duration | 6 WEEKS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

LUMACAFTOR-IVACAFTOR

Products Affected

• ORKAMBI ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF. |
| Age Restrictions | |
| Prescriber Restrictions | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: LIFETIME. |
| Other Criteria | CF: RENEWAL: IMPROVEMENT IN CLINICAL STATUS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MACITENTAN

Products Affected

OPSUMIT

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MARGETUXIMAB-CMKB

Products Affected

MARGENZA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MARIBAVIR

Products Affected

LIVTENCITY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MECASERMIN

Products Affected

INCRELEX

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR NEPHROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF WRIST AND HAND. RENEWAL: IMPROVEMENT WHILE ON THERAPY (I.E., INCREASE IN HEIGHT OR INCREASE IN HEIGHT VELOCITY). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MECHLORETHAMINE

Products Affected

• VALCHLOR

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MEPOLIZUMAB

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. |
| Coverage Duration | INITIAL: ASTHMA: 12 MO. CRSWNP: 6 MO. OTHERS: 12 MO. RENEWAL: CRSWNP, ASTHMA: 12 MO. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-ILS BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. RENEWAL: ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-ILS BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE. CRSWNP: 1) CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MIDOSTAURIN

Products Affected

RYDAPT

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MIFEPRISTONE

Products Affected

• mifepristone oral tablet 300 mg

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM). |
| Age Restrictions | |
| Prescriber Restrictions | CS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS |
| Other Criteria | CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO THERAPY, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MILTEFOSINE

Products Affected

IMPAVIDO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MIRDAMETINIB

- GOMEKLI ORAL CAPSULE 1 MG, 2 MG
- GOMEKLI ORAL TABLET FOR SUSPENSION

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MIRVETUXIMAB SORAVTANSINE-GYNX

Products Affected

• ELAHERE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: AN OPHTHALMIC EXAM, INCLUDING VISUAL ACUITY AND SLIT LAMP EXAM, WILL BE COMPLETED PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MOMELOTINIB

Products Affected

OJJAARA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

MOSUNETUZUMAB-AXGB

Products Affected

• LUNSUMIO

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS. |
| Other Criteria | RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NARCOLEPSY AGENTS

- armodafinil
- modafinil oral tablet 100 mg, 200 mg

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NAXITAMAB-GQGK

Products Affected

DANYELZA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NERATINIB

Products Affected

NERLYNX

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NILOTINIB

Products Affected

• TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NILOTINIB-DANZITEN

Products Affected

DANZITEN

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): 1) PERFORMED MUTATIONAL ANALYSIS PRIOR TO INITIATION OF THERAPY, AND 2) THERAPY IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NINTEDANIB

Products Affected

• OFEV

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENED/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIRAPARIB

Products Affected

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIRAPARIB-ABIRATERONE

Products Affected

AKEEGA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIROGACESTAT

Products Affected

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NITISINONE

Products Affected

- nitisinone
- ORFADIN ORAL SUSPENSION

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE. |
| Age Restrictions | |
| Prescriber Restrictions | HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. |
| Other Criteria | HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIVOLUMAB

Products Affected

OPDIVO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIVOLUMAB-HYALURONIDASE-NVHY

Products Affected

• OPDIVO QVANTIG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NIVOLUMAB-RELATLIMAB-RMBW

Products Affected

OPDUALAG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

NOGAPENDEKIN ALFA

Products Affected

ANKTIVA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 40 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OCRELIZUMAB

Products Affected

OCREVUS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OCRELIZUMAB-HYALURONIDASE-OCSQ

Products Affected

OCREVUS ZUNOVO

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OFATUMUMAB-SQ

Products Affected

KESIMPTA PEN

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OLANZAPINE/SAMIDORPHAN

Products Affected

LYBALVI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | SCHIZOPHRENIA, BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST |
| Coverage Duration | 12 MONTHS |
| Other Criteria | SCHIZOPHRENIA: 1) AT HIGH RISK FOR WEIGHT GAIN, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF LURASIDONE OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: 1) AT HIGH RISK FOR WEIGHT GAIN, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OLAPARIB

Products Affected

LYNPARZA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: MEDICATION WILL BE USED AS MONOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OLUTASIDENIB

Products Affected

REZLIDHIA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OMACETAXINE

Products Affected

SYNRIBO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OMALIZUMAB

Products Affected

XOLAIR

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL OF AT LEAST 30 IU/ML. FOOD ALLERGY: 1) IGE SERUM LEVEL OF AT LEAST 30 IU/ML, AND 2) ALLERGEN SPECIFIC IGE SERUM LEVEL OF AT LEAST 6 KUA/L TO AT LEAST ONE FOOD, OR POSITIVE SKIN PRICK TEST TO AT LEAST ONE FOOD, OR POSITIVE MEDICALLY MONITORED FOOD CHALLENGE TO AT LEAST ONE FOOD. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL/RENEWAL: CHRONIC SPONTANEOUS URTICARIA (CSU): PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, DERMATOLOGIST, OR IMMUNOLOGIST. INITIAL: CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. FOOD ALLERGY: PRESCRIBED BY OR IN CONSULTATION WITH ALLERGIST OR IMMUNOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: ASTHMA 12 MO/12 MO, CSU 6 MO/12 MO, CRSWNP 6 MO/12 MO, FOOD ALLERGY 12 MO/24 MO |

| | , |
|----------------------------|---|
| PA Criteria | Criteria Details |
| PA Criteria Other Criteria | INITIAL: CSU: 1) TRIAL OF AND MAINTAINED ON, OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE AND 2) STILL EXPERIENCES HIVES OR ANGIOEDEMA ON MOST DAYS OF THE WEEK FOR AT LEAST 6 WEEKS. CRSWNP: 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT, AND 3) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH DUPIXENT, TEZSPIRE, OR ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. FOOD ALLERGY: 1) |
| | ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH DUPIXENT, TEZSPIRE, OR ANTI-IL5 |
| | EPINEPHRINE AUTO-INJECTOR/INJECTION, AND 2) NO CONCURRENT USE WITH PEANUT-SPECIFIC IMMUNOTHERAPY. RENEWAL: CSU: MAINTAINED ON OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI- |
| | HISTAMINE. CRSWNP: 1) CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., |

| PA Criteria | Criteria Details |
|------------------------|--|
| | JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. ASTHMA: 1) NO CONCURRENT USE WITH DUPIXENT, TEZSPIRE, OR ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMARELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. FOOD ALLERGY: 1) PERSISTENT IGE-MEDIATED FOOD ALLERGY, 2) CONCURRENT USE WITH AN ACTIVE PRESCRIPTION FOR EPINEPHRINE AUTO-INJECTOR/INJECTION, AND 3) NO CONCURRENT USE WITH PEANUT-SPECIFIC IMMUNOTHERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OSIMERTINIB

Products Affected

TAGRISSO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

OXANDROLONE

Products Affected

• oxandrolone

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 6 MONTHS |
| Other Criteria | PROTEIN CATABOLISM, BONE PAIN: 1) MONITORED FOR PELIOSIS HEPATIS, LIVER CELL TUMORS, AND BLOOD LIPID CHANGES, 2) DOES NOT HAVE KNOWN OR SUSPECTED: CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, NEPHROSIS (THE NEPHROTIC PHASE OF NEPHRITIS), OR HYPERCALCEMIA, AND 3) DOES NOT HAVE SEVERE HEPATIC DYSFUNCTION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PACRITINIB

Products Affected

VONJO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PALBOCICLIB

Products Affected

• IBRANCE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE PREFERRED AGENTS, WHERE INDICATIONS ALIGN: KISQALI, VERZENIO. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PARATHYROID HORMONE

Products Affected

NATPARA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PASIREOTIDE DIASPARTATE

Products Affected

SIGNIFOR

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. |
| Other Criteria | CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PAZOPANIB

Products Affected

pazopanib

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST) |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEGFILGRASTIM - APGF

Products Affected

NYVEPRIA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEGFILGRASTIM-NEULASTA ONPRO

Products Affected

NEULASTA ONPRO

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEGINTERFERON ALFA-2A

Products Affected

PEGASYS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | HEPATITIS B: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, OR PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G., HEPATOLOGIST). |
| Coverage Duration | HEP B/HEP C: 48 WEEKS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEGVISOMANT

Products Affected

SOMAVERT

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEMBROLIZUMAB

Products Affected

KEYTRUDA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEMIGATINIB

Products Affected

PEMAZYRE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | CHOLANGIOCARCINOMA, MYELOID/LYMPHOID NEOPLASMS: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PENICILLAMINE TABLET

Products Affected

• penicillamine oral tablet

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: CYSTINURIA: HAS NEPHROLITHIASIS AND ONE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTINE, 2) PRESENCE OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) FAMILY HISTORY OF CYSTINURIA AND POSITIVE CYANIDE-NITROPRUSSIDE SCREENING. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: 12 MONTHS, RENEWAL: LIFETIME. |
| Other Criteria | INITIAL: WILSONS DISEASE: 1) LEIPZIG SCORE OF 4 OR GREATER. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: CONTINUES TO BENEFIT FROM THE MEDICATION. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PEXIDARTINIB

Products Affected

TURALIO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PIMAVANSERIN

Products Affected

NUPLAZID

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER |
| Prescriber Restrictions | PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST). |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PIRFENIDONE

Products Affected

- pirfenidone oral capsulepirfenidone oral tablet 267 mg, 534 mg, 801

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE. |
| Age Restrictions | IPF: INITIAL: 18 YEARS OR OLDER. |
| Prescriber Restrictions | IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PIRTOBRUTINIB

Products Affected

 JAYPIRCA ORAL TABLET 100 MG, 50 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

POMALIDOMIDE

Products Affected

POMALYST

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PONATINIB

Products Affected

• ICLUSIG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | CHRONIC MYELOID LEUKEMIA (CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

POSACONAZOLE TABLET

Products Affected

• posaconazole oral tablet,delayed release (dr/ec)

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE, PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PRALSETINIB

Products Affected

GAVRETO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

PYRIMETHAMINE

Products Affected

• pyrimethamine

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST. |
| Coverage Duration | TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS. |
| Other Criteria | TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

QUININE

Products Affected

• quinine sulfate

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

QUIZARTINIB

Products Affected

VANFLYTA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

REGORAFENIB

Products Affected

STIVARGA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RELUGOLIX

Products Affected

ORGOVYX

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

REPOTRECTINIB

Products Affected

AUGTYRO ORAL CAPSULE 160 MG, 40 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RESLIZUMAB

Products Affected

CINQAIR

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. |
| Age Restrictions | |
| Prescriber Restrictions | ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RETIFANLIMAB-DLWR

Products Affected

ZYNYZ

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

REVUMENIB

Products Affected

 REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RIBOCICLIB

Products Affected

 KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RIBOCICLIB-LETROZOLE

Products Affected

 KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RIFAXIMIN

Products Affected

• XIFAXAN ORAL TABLET 200 MG, 550 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | TRAVELERS DIARRHEA, HEPATIC ENCEPHALOPATHY (HE): 12 MOS. IBS-D: 8 WKS. |
| Other Criteria | HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RILONACEPT

Products Affected

ARCALYST

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF- FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR \$100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES. DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS. RECURRENT PERICARDITIS (RP): TWO OF THE FOLLOWING: CHEST PAIN CONSISTENT WITH PERICARDITIS, PERICARDIAL FRICTION RUB, ECG SHOWING DIFFUSE ST-SEGMENT ELEVATION OR PR- SEGMENT DEPRESSION, NEW OR WORSENING PERICARDIAL EFFUSION. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CAPS, DIRA: LIFETIME. RP: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | CAPS: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. DIRA: 1) NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS, AND 2) TRIAL OF THE PREFERRED AGENT: KINERET. RP: 1) HAD AN EPISODE OF ACUTE PERICARDITIS, 2) SYMPTOM-FREE FOR 4 TO 6 WEEKS, AND 3) NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RIMEGEPANT

Products Affected

NURTEC ODT

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: ACUTE MIGRAINE TREATMENT: 1) TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN), AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT, AND 2) ONE OF THE FOLLOWING: (A) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR (B) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RIOCIGUAT

Products Affected

ADEMPAS

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) (WHO GROUP 4): WHO FUNCTIONAL CLASS II-IV SYMPTOMS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: PAH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE (PDE) INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

RIPRETINIB

Products Affected

QINLOCK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RISANKIZUMAB-RZAA

Products Affected

SKYRIZI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSO. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RITUXIMAB AND HYALURONIDASE HUMAN-SQ

Products Affected

RITUXAN HYCELA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | FOLLICULAR LYMPHOMA (FL), DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RITUXIMAB-ABBS

Products Affected

TRUXIMA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST. |
| Coverage Duration | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO. |
| Other Criteria | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RITUXIMAB-ARRX

Products Affected

RIABNI

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST. |
| Coverage Duration | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO. |
| Other Criteria | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RITUXIMAB-PVVR

Products Affected

RUXIENCE

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | RA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST. |
| Coverage Duration | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO. |
| Other Criteria | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ROPEGINTERFERON ALFA-2B-NJFT

Products Affected

BESREMI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RUCAPARIB

Products Affected

RUBRACA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

RUXOLITINIB

Products Affected

JAKAFI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS |
| Other Criteria | MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SAPROPTERIN

Products Affected

- javygtor oral tablet, soluble
- sapropterin oral tablet, soluble

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 2 MONTHS, RENEWAL 12 MONTHS. |
| Other Criteria | HYPERPHENYLALANINEMIA (HPA): INITIAL: NO CONCURRENT USE WITH PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NO CONCURRENT USE WITH PALYNZIQ. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SECUKINUMAB IV

Products Affected

COSENTYX INTRAVENOUS

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI). |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SECUKINUMAB SQ

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML

• COSENTYX UNOREADY PEN

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI). |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: HS: 12 MONTHS, ALL OTHER INDICATIONS: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|--|
| Other Criteria | INITIAL: PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSO. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NS. AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. RENEWAL: PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |
| | MOLECULES FOR NR-AXSPA. ERA, HS: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

SELEXIPAG

Products Affected

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS, DOSE PACK

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS |
| Other Criteria | PAH: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SELINEXOR

Products Affected

• XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (10 MG X 4), 40 MG/WEEK (20 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SELPERCATINIB

Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG
- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SELUMETINIB

Products Affected

• KOSELUGO ORAL CAPSULE 10 MG, 25 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SILDENAFIL TABLET

Products Affected

• sildenafil (pulm.hypertension) oral tablet

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: AGES 18 YEARS OR OLDER: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. AGES 1 TO 17 YEARS: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PAP GREATER THAN 20 MMHG, 2) PCWP OF 15 MMHG OR LESS, AND 3) PVR OF 3 WOOD UNITS OR GREATER. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SIPONIMOD

Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER (FOR 2MG MAINT)

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SIROLIMUS PROTEIN-BOUND

Products Affected

FYARRO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SODIUM OXYBATE-XYREM

Products Affected

• sodium oxybate

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: CATAPLEXY IN NARCOLEPSY, EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS |
| Other Criteria | INITIAL: EDS IN NARCOLEPSY: 1) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT, 2) AGES 18 YEARS OR OLDER: TRIAL, FAILURE OF, OR CONTRAINDICATION TO A FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, OR SUNOSI AND ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) AGES 7 TO 17 YEARS: TRIAL, FAILURE OF, OR CONTRAINDICATION TO ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT. RENEWAL: CATAPLEXY IN NARCOLEPSY, EDS IN NARCOLEPSY: 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOFOSBUVIR/VELPATASVIR

Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. |
| Other Criteria | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANAVIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

Products Affected

VOSEVI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE. |
| Other Criteria | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOMATROPIN - NORDITROPIN

Products Affected

NORDITROPIN FLEXPRO

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES. |
| Required Medical Information | INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS BELOW THE MEAN HEIGHT FOR CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL/RENEWAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | INITIAL: ADULT GHD: GHD ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASE, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, OR TRAUMA, OR FOR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GHD. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR HAS NOT COMPLETED PREPUBERTAL GROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION. |

| PA Criteria | Criteria Details |
|------------------------|-------------------------------|
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOMATROPIN - SEROSTIM

Products Affected

• SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES |
| Required Medical Information | INITIAL: HIV/WASTING: ONE OF THE FOLLOWING FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BCM LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND BMI LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED. |
| Age Restrictions | |
| Prescriber Restrictions | HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST. |
| Coverage Duration | INITIAL/RENEWAL: 3 MONTHS. |
| Other Criteria | HIV/WASTING: INITIAL: 1) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SONIDEGIB

Products Affected

ODOMZO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC): BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SORAFENIB

Products Affected

sorafenib

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOTATERCEPT-CSRK

Products Affected

WINREVAIR

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | PAH: INITIAL: 1) ON BACKGROUND PAH THERAPY (FOR AT LEAST 3 MONTHS) WITH AT LEAST TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: A) ORAL ENDOTHELIN RECEPTOR ANTAGONIST, B) ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, C) ORAL CGMP STIMULATOR, D) IV/SQ PROSTACYCLIN, OR 2) ON ONE AGENT FROM ONE OF THE ABOVE DRUG CLASSES, AND HAS A CONTRAINDICATION OR INTOLERANCE TO ALL OF THE OTHER DRUG CLASSES. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SOTORASIB

Products Affected

• LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

STIRIPENTOL

Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

SUNITINIB

Products Affected

• sunitinib malate

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC). |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TADALAFIL - ADCIRCA, ALYQ

Products Affected

alyq

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions | |
| Prescriber Restrictions | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS. |
| Other Criteria | PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM, AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS. |
| Indications | All Medically-accepted Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TADALAFIL-CIALIS

Products Affected

• tadalafil oral tablet 2.5 mg, 5 mg

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH). |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | BPH: 1) TRIAL OF ONE ALPHA BLOCKER (E.G., DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (E.G., FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TALAZOPARIB

Products Affected

TALZENNA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | ADVANCED OR METASTATIC BREAST CANCER: 1) HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING, AND 2) IF HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER, RECEIVED PRIOR TREATMENT WITH ENDOCRINE THERAPY OR IS CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TALQUETAMAB-TGVS

Products Affected

TALVEY

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TARLATAMAB-DLLE

Products Affected

IMDELLTRA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TAZEMETOSTAT

Products Affected

TAZVERIK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TEBENTAFUSP-TEBN

Products Affected

KIMMTRAK

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TECLISTAMAB-CQYV

Products Affected

TECVAYLI

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TELOTRISTAT

Products Affected

XERMELO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TEPOTINIB

Products Affected

TEPMETKO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TERIPARATIDE

Products Affected

• teriparatide subcutaneous pen injector 20 mcg/dose (620mcg/2.48ml)

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 24 MONTHS |
| Other Criteria | OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY, UNLESS REMAINS AT OR HAS RETURNED TO HAVING A HIGH RISK FOR FRACTURE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TESTOSTERONE

- testosterone transdermal gel in metered-dose pump 12.5 mg/ 1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)
- $(25 \ mg/2.5 gram), \ 1 \% \ (50 \ mg/5 \ gram)$

| • | testosterone | transdermal | gel i | in packet | 1 | $\frac{9}{6}$ | ó |
|---|--------------|-------------|-------|-----------|---|---------------|---|
|---|--------------|-------------|-------|-----------|---|---------------|---|

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS |
| Other Criteria | MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TESTOSTERONE CYPIONATE

Products Affected

• testosterone cypionate

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL: 12 MONTHS |
| Other Criteria | MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TESTOSTERONE ENANTHATE

- testosterone enanthate
- XYOSTED

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL. |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL: MALE DELAYED PUBERTY: 6MO, MALE HYPOGONADISM: 12 MO. OTHER INDICATIONS: 12 MO. |
| Other Criteria | INITIAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. MALE DELAYED PUBERTY: HAS NOT RECEIVED MORE THAN TWO 6-MONTH COURSES OF TESTOSTERONE REPLACEMENT THERAPY |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TETRABENAZINE

Products Affected

tetrabenazine

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

THALIDOMIDE

Products Affected

THALOMID

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TISLELIZUMAB-JSGR

Products Affected

• TEVIMBRA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TISOTUMAB VEDOTIN-TFTV

Products Affected

• TIVDAK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TIVOZANIB

Products Affected

FOTIVDA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOCILIZUMAB IV

Products Affected

ACTEMRA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. |
| Coverage Duration | INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, RINVOQ, ORENCIA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOCILIZUMAB SQ

- ACTEMRA
- ACTEMRA ACTPEN

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, RINVOQ, ORENCIA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOCILIZUMAB-AAZG

- TYENNE
- TYENNE AUTOINJECTOR

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOFACITINIB

- XELJANZ
- XELJANZ XR

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PCJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PCJIA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PCJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PCJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PCJIA. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PCJIA. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOPICAL TRETINOIN

- ALTRENO
- tretinoin topical cream

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA. |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TORIPALIMAB-TPZI

Products Affected

• LOQTORZI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | NASOPHARYNGEAL CARCINOMA (NPC): FIRST LINE TREATMENT: 24 MOS, PREVIOUSLY TREATED: LIFETIME. |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TOVORAFENIB

- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION
- OJEMDA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRAMETINIB SOLUTION

Products Affected

• MEKINIST ORAL RECON SOLN

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | UNRESECTABLE OR METASTATIC MELANOMA, MELANOMA, METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), LOCALLY ADVANCED OR METASTATIC ANAPLASTIC THYROID CANCER (ATC), UNRESECTABLE OR METASTATIC SOLID TUMOR, LOW-GRADE GLIOMA (LGG): UNABLE TO SWALLOW MEKINIST TABLETS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRAMETINIB TABLET

Products Affected

• MEKINIST ORAL TABLET 0.5 MG, 2 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRASTUZUMAB-DKST

Products Affected

OGIVRI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRASTUZUMAB-DTTB

Products Affected

ONTRUZANT

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRASTUZUMAB-HYALURONIDASE-OYSK

Products Affected

• HERCEPTIN HYLECTA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRASTUZUMAB-PKRB

Products Affected

HERZUMA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRASTUZUMAB-QYYP

Products Affected

TRAZIMERA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRAZODONE

Products Affected

RALDESY

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | MAJOR DEPRESSIVE DISORDER (MDD): CONTRAINDICATION TO OR UNABLE TO SWALLOW TRAZODONE TABLETS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TREMELIMUMAB-ACTL

Products Affected

• IMJUDO

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS. |
| Other Criteria | UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO. NSCLC: HAS NOT RECEIVED A TOTAL OF 5 DOSES OF IMJUDO. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRIENTINE CAPSULE

Products Affected

• trientine oral capsule 250 mg

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 12 MONTHS, RENEWAL: LIFETIME. |
| Other Criteria | WILSONS DISEASE: INITIAL: 1) LEIPZIG SCORE OF 4 OR GREATER, AND 2) TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE TABLET. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRIFLURIDINE/TIPIRACIL

Products Affected

• LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TRIPTORELIN-TRELSTAR

Products Affected

• TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS. |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

TUCATINIB

Products Affected

• TUKYSA ORAL TABLET 150 MG, 50 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

UBROGEPANT

Products Affected

UBRELVY

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |
| Other Criteria | ACUTE MIGRAINE TREATMENT: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN), AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. RENEWAL: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT, AND 2) ONE OF THE FOLLOWING: (A) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR (B) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

UPADACITINIB

- RINVOQRINVOQ LQ

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI). ATOPIC DERMATITIS (AD): ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. ULCERATIVE COLITIS (UC), CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|----------------|---|
| Other Criteria | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 2) TRIAL OF OR CONTRAINDICATION TO A TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITORS FOR ANY INDICATION. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG), AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITOR FOR ANY INDICATION. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITOR FOR ANY INDICATION. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITOR FOR ANY INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITOR FOR ANY INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC |

| PA Criteria | Criteria Details |
|------------------------|--|
| | CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. PJIA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. TARGETED SMALL MOLECULES FOR CD. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB

Products Affected

• STELARA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB IV

Products Affected

• STELARA

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | 2 MONTHS |
| Other Criteria | CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB-AEKN IV

Products Affected

SELARSDI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB-AEKN SQ

Products Affected

SELARSDI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB-KFCE IV

Products Affected

YESINTEK

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|---|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

USTEKINUMAB-KFCE SQ

Products Affected

YESINTEK

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE. |
| Age Restrictions | |
| Prescriber Restrictions | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS. |

| PA Criteria | Criteria Details |
|------------------------|--|
| Other Criteria | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSO. 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSO. 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VALBENAZINE

Products Affected

- INGREZZA
- INGREZZA INITIATION PK(TARDIV)
- INGREZZA SPRINKLE

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST. CHOREA ASSOCIATED WITH HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | TD: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VANDETANIB

Products Affected

CAPRELSA ORAL TABLET 100 MG, 300 MG

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VANZACAFTOR-TEZACAFTOR-DEUTIVACAFTOR

Products Affected

• ALYFTREK ORAL TABLET 10-50-125 MG, 4-20-50 MG

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT. |
| Coverage Duration | INITIAL: 6 MONTHS. RENEWAL: LIFETIME. |
| Other Criteria | CF: INITIAL: NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VEMURAFENIB

Products Affected

ZELBORAF

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VENETOCLAX

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VERICIGUAT

Products Affected

VERQUVO

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | INITIAL/RENEWAL:12 MONTHS. |
| Other Criteria | HEART FAILURE (HF): INITIAL: 1) NO CONCURRENT USE WITH LONG-ACTING NITRATES OR NITRIC OXIDE DONORS, RIOCIGUAT, OR PDE-5 INHIBITORS, 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED SGLT-2 INHIBITOR, AND 3) TRIAL OF OR CONTRAINDICATION TO ONE AGENT FROM ANY OF THE FOLLOWING STANDARD OF CARE CLASSES: (A) ACE INHIBITOR, ARB, OR ARNI, (B) BETA BLOCKER (I.E., BISOPROLOL, CARVEDILOL, METOPROLOL SUCCINATE), OR (C) ALDOSTERONE ANTAGONIST (I.E., SPIRONOLACTONE, EPLERENONE). RENEWAL: NO CONCURRENT USE WITH LONG-ACTING NITRATES OR NITRIC OXIDE DONORS, RIOCIGUAT, OR PDE-5 INHIBITORS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VIGABATRIN

Products Affected

- vigabatrin
- vigadronevigpoder

| PA Criteria | Criteria Details |
|------------------------------------|---|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage Duration | 12 MONTHS |
| Other Criteria | CPS: TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VIMSELTINIB

Products Affected

ROMVIMZA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VISMODEGIB

Products Affected

• ERIVEDGE

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VORASIDENIB

Products Affected

VORANIGO

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

VORICONAZOLE SUSPENSION

Products Affected

• voriconazole oral suspension for reconstitution

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | CANDIDA INFECTIONS: 3 MOS. CONTINUATION OF THERAPY, ALL OTHER INDICATIONS: 6 MOS. |
| Other Criteria | CANDIDA INFECTIONS: 1) TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE, AND 2) UNABLE TO SWALLOW TABLETS. ALL INDICATIONS EXCEPT ESOPHAGEAL CANDIDIASIS: UNABLE TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ZANIDATAMAB-HRII

Products Affected

ZIIHERA

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ZANUBRUTINIB

Products Affected

BRUKINSA

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ZENOCUTUZUMAB-ZBCO

Products Affected

BIZENGRI

| PA Criteria | Criteria Details |
|------------------------------------|--|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ZOLBETUXIMAB-CLZB

Products Affected

VYLOY

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 12 MONTHS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

ZURANOLONE

Products Affected

• ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

| PA Criteria | Criteria Details |
|------------------------------------|-------------------------------|
| Exclusion Criteria | |
| Required Medical Information | |
| Age Restrictions | |
| Prescriber Restrictions | |
| Coverage Duration | 14 DAYS |
| Other Criteria | |
| Indications | All FDA-approved Indications. |
| Off Label Uses | |
| Part B Prerequisite | No |

INDEX

| INDEA | |
|-------------------------------------|-------------------------------------|
| 1ST TIER UNIFINE PENTP 5MM 31G 157 | ALUNBRIG ORAL TABLETS,DOSE |
| 1ST TIER UNIFINE PNTIP 4MM 32G157 | PACK |
| 1ST TIER UNIFINE PNTIP 6MM 31G157 | ALVAIZ104 |
| 1ST TIER UNIFINE PNTIP 8MM 31G | ALYFTREK ORAL TABLET 10-50-125 |
| STRL,SINGLE-USE,SHRT157 | MG, 4-20-50 MG374 |
| 1ST TIER UNIFINE PNTP 29GX1/2" 157 | <i>alyq</i> |
| 1ST TIER UNIFINE PNTP 31GX3/16157 | ANKTIVA226 |
| 1ST TIER UNIFINE PNTP 32GX5/32157 | AQINJECT PEN NEEDLE 31G 5MM157 |
| abiraterone7 | AQINJECT PEN NEEDLE 32G 4MM 157 |
| abirtega7 | ARCALYST273 |
| ABOUTTIME PEN NEEDLE157 | ARIKAYCE23 |
| ACTEMRA333, 335 | armodafinil212 |
| ACTEMRA ACTPEN335 | ASSURE ID DUO PRO NDL 31G 5MM157 |
| ACTHAR75 | ASSURE ID DUO-SHIELD 30GX3/16"157 |
| ACTHAR SELFJECT SUBCUTANEOUS | ASSURE ID DUO-SHIELD 30GX5/16"157 |
| PEN INJECTOR 40 UNIT/0.5 ML, 80 | ASSURE ID INSULIN SAFETY |
| UNIT/ML75 | SYRINGE 1 ML 29 GAUGE X 1/2" 157 |
| ACTIMMUNE170 | ASSURE ID PEN NEEDLE 30GX3/16"157 |
| ADEMPAS277 | ASSURE ID PEN NEEDLE 30GX5/16"157 |
| ADVOCATE INS 0.3 ML 30GX5/16" 157 | ASSURE ID PEN NEEDLE 31GX3/16"157 |
| ADVOCATE INS 0.3 ML 31GX5/16" 157 | ASSURE ID PRO PEN NDL 30G 5MM157 |
| ADVOCATE INS 0.5 ML 30GX5/16" 157 | ASSURE ID SYR 0.5 ML 29GX1/2" (RX) |
| ADVOCATE INS 0.5 ML 31GX5/16" 157 | 157 |
| ADVOCATE INS 1 ML 31GX5/16" 157 | ASSURE ID SYR 0.5 ML 31GX15/64" 157 |
| ADVOCATE INS SYR 0.3 ML 29GX1/2.157 | ASSURE ID SYR 1 ML 31GX15/64" 157 |
| ADVOCATE INS SYR 0.5 ML 29GX1/2.157 | AUGTYRO ORAL CAPSULE 160 MG, |
| ADVOCATE INS SYR 1 ML 29GX1/2" 157 | 40 MG265 |
| ADVOCATE INS SYR 1 ML 30GX5/16 157 | AUSTEDO ORAL TABLET 12 MG, 6 |
| ADVOCATE PEN NDL 12.7MM 29G 157 | MG, 9 MG |
| ADVOCATE PEN NEEDLE 32G 4MM 157 | AUSTEDO XR ORAL TABLET |
| ADVOCATE PEN NEEDLE 4MM 33G 157 | EXTENDED RELEASE 24 HR 12 MG, 18 |
| ADVOCATE PEN NEEDLES 5MM 31G.157 | MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 |
| ADVOCATE PEN NEEDLES 8MM 31G.157 | MG, 6 MG |
| AJOVY AUTOINJECTOR131 | AUSTEDO XR TITRATION KT(WK1-4)89 |
| AJOVY SYRINGE131 | AUTOSHIELD DUO PEN NDL 30G |
| AKEEGA220 | 5MM |
| ALCOHOL 70% SWABS157 | AVONEX INTRAMUSCULAR PEN |
| ALCOHOL PADS157 | INJECTOR KIT167 |
| ALCOHOL PREP SWABS157 | AVONEX INTRAMUSCULAR |
| ALCOHOL WIPES157 | SYRINGE KIT |
| ALECENSA | AVONEX PEN 30 MCG/0.5 ML167 |
| ALTRENO341 | AYVAKIT |
| ALUNBRIG ORAL TABLET 180 MG, 30 | BALVERSA ORAL TABLET 3 MG, 4 |
| MG, 90 MG | MG, 5 MG |
| 1120, 20 1110 | 117 |

| BD AUTOSHIELD DUO NDL | BENDAMUSTINE INTRAVENOUS |
|--|------------------------------------|
| 5MMX30G157 | SOLUTION48 |
| BD ECLIPSE 30GX1/2" SYRINGE157 | BENDEKA48 |
| BD ECLIPSE NEEDLE 30GX1/2" (OTC) 157 | BENLYSTA SUBCUTANEOUS45 |
| BD INS SYR 0.3 ML 8MMX31G(1/2)157 | BESREMI286 |
| BD INS SYR UF 0.3 ML 12.7MMX30G 157 | <i>betaine</i> 51 |
| BD INS SYR UF 0.5 ML 12.7MMX30G | BETASERON SUBCUTANEOUS KIT 168 |
| NOT FOR RETAIL SALE157 | <i>bexarotene</i> 55 |
| BD INS SYRN UF 1 ML 12.7MMX30G | BIZENGRI385 |
| NOT FOR RETAIL SALE157 | BORDERED GAUZE 2"X2"157 |
| BD INS SYRNG UF 0.3 ML 8MMX31G157 | bortezomib injection57 |
| BD INS SYRNG UF 0.5 ML 8MMX31G157 | BORUZU57 |
| BD INSULIN SYR 1 ML 25GX1" 157 | bosentan58 |
| BD INSULIN SYR 1 ML 25GX5/8" 157 | BOSULIF ORAL CAPSULE 100 MG, 50 |
| BD INSULIN SYR 1 ML 26GX1/2" 157 | MG59 |
| BD INSULIN SYR 1 ML 27GX12.7MM157 | BOSULIF ORAL TABLET 100 MG, 400 |
| BD INSULIN SYR 1 ML 27GX5/8" | MG, 500 MG 59 |
| BD INSULIN SYR 1 ML 27GX5/8" MICRO-FINE157 | BRAFTOVI107 |
| BD INSULIN SYRINGE SLIP TIP157 | BRUKINSA384 |
| BD INSULIN SYRINGE U-500157 | CABOMETYX ORAL TABLET 20 MG, |
| BD LUER-LOK SYRINGE 1 ML 157 | 40 MG, 60 MG63 |
| BD NANO 2 GEN PEN NDL 32G 4MM 157 | CALQUENCE9 |
| BD SAFETGLD INS 0.3 ML 29G 13MM.157 | CALQUENCE (ACALABRUTINIB |
| BD SAFETGLD INS 0.5 ML 13MMX29G | MAL)9 |
| | CAPRELSA ORAL TABLET 100 MG, |
| BD SAFETYGLD INS 0.3 ML 31G 8MM 157 | 300 MG373 |
| BD SAFETYGLD INS 0.5 ML 30G 8MM 157 | CAREFINE PEN NEEDLE 12.7MM 29G.157 |
| BD SAFETYGLD INS 1 ML 29G 13MM.157 | CAREFINE PEN NEEDLE 4MM 32G157 |
| BD SAFETYGLID INS 1 ML 6MMX31G157 | CAREFINE PEN NEEDLE 5MM 32G157 |
| BD SAFETYGLIDE SYRINGE 27GX5/8 157 | CAREFINE PEN NEEDLE 6MM 31G157 |
| BD SAFTYGLD INS 0.3 ML 6MMX31G 157 | CAREFINE PEN NEEDLE 8MM 30G157 |
| BD SAFTYGLD INS 0.5 ML 29G 13MM 157 | CAREFINE PEN NEEDLES 6MM 32G 157 |
| BD SAFTYGLD INS 0.5 ML 6MMX31G 157 | CAREFINE PEN NEEDLES 8MM 31G 157 |
| BD SINGLE USE SWAB157 | CARETOUCH ALCOHOL 70% PREP |
| BD UF MICRO PEN NEEDLE | PAD |
| 6MMX32G | CARETOUCH PEN NEEDLE 29G 12MM |
| BD UF MINI PEN NEEDLE 5MMX31G. 157 | |
| BD UF NANO PEN NEEDLE 4MMX32G | CARETOUCH PEN NEEDLE 31GX1/4". 157 |
| | CARETOUCH PEN NEEDLE 31GX3/16" |
| BD UF ORIG PEN NDL 12.7MMX29G 157 | |
| BD UF SHORT PEN NEEDLE | CARETOUCH PEN NEEDLE 31GX5/16" |
| 8MMX31G | |
| BD VEO INS 0.3 ML 6MMX31G (1/2) 157 | CARETOUCH PEN NEEDLE 32GX3/16" |
| BD VEO INS SYRING 1 ML 6MMX31G 157 | |
| BD VEO INS SYRN 0.3 ML 6MMX31G. 157 | CARETOUCH PEN NEEDLE 32GX5/32" |
| BD VEO INS SYRN 0.5 ML 6MMX31G. 157 | |
| bendamustine intravenous recon soln 48 | CARETOUCH SYR 0.3 ML 31GX5/16" 157 |

| CARETOUCH SYR 0.5 ML 30GX5/16" 157 | COMFORT EZ PEN NEEDLES 6MM |
|------------------------------------|------------------------------------|
| CARETOUCH SYR 0.5 ML 31GX5/16" 157 | 33G157 |
| CARETOUCH SYR 1 ML 28GX5/16" 157 | COMFORT EZ PEN NEEDLES 8MM |
| CARETOUCH SYR 1 ML 29GX5/16" 157 | 31G SHORT157 |
| CARETOUCH SYR 1 ML 30GX5/16" 157 | COMFORT EZ PEN NEEDLES 8MM |
| CARETOUCH SYR 1 ML 31GX5/16" 157 | 32G157 |
| carglumic acid67 | COMFORT EZ PEN NEEDLES 8MM |
| CAYSTON43 | 33G157 |
| CIMZIA POWDER FOR RECONST69 | COMFORT EZ PRO PEN NDL 30G 8MM |
| CIMZIA SUBCUTANEOUS SYRINGE | |
| KIT 400 MG/2 ML (200 MG/ML X 2)69 | COMFORT EZ PRO PEN NDL 31G 4MM |
| CINQAIR266 | |
| CLICKFINE 31G X 5/16" NEEDLES | COMFORT EZ PRO PEN NDL 31G 5MM |
| 8MM, UNIVERSAL 157 | |
| CLICKFINE PEN NEEDLE 32GX5/32" | COMFORT EZ SYR 0.3 ML 29GX1/2"157 |
| 32GX4MM, STERILE157 | COMFORT EZ SYR 0.5 ML 28GX1/2"157 |
| CLICKFINE UNIVERSAL 31G X 1/4" | COMFORT EZ SYR 0.5 ML 29GX1/2"157 |
| 6MM, STORE BRAND 157 | COMFORT EZ SYR 0.5 ML 30GX1/2"157 |
| COMETRIQ ORAL CAPSULE 100 | COMFORT EZ SYR 1 ML 28GX1/2"157 |
| MG/DAY(80 MG X1-20 MG X1), 140 | COMFORT EZ SYR 1 ML 29GX1/2"157 |
| MG/DAY(80 MG X1-20 MG X3), 60 | COMFORT EZ SYR 1 ML 30GX1/2"157 |
| MG/DAY (20 MG X 3/DAY)62 | COMFORT EZ SYR 1 ML 30GX5/16"157 |
| COMFORT EZ 0.3 ML 31G 15/64" 157 | COMFORT POINT PEN NDL 31GX1/3".157 |
| COMFORT EZ 0.5 ML 31G 15/64" 157 | COMFORT POINT PEN NDL 31GX1/6".157 |
| COMFORT EZ INS 0.3 ML 30GX1/2"157 | COMFORT TOUCH PEN NDL 31G 4MM |
| COMFORT EZ INS 0.3 ML 30GX5/16"157 | |
| COMFORT EZ INS 1 ML 31G 15/64"157 | COMFORT TOUCH PEN NDL 31G 5MM |
| COMFORT EZ INS 1 ML 31GX5/16"157 | |
| COMFORT EZ INSULIN SYR 0.3 ML 157 | COMFORT TOUCH PEN NDL 31G 6MM |
| COMFORT EZ INSULIN SYR 0.5 ML 157 | |
| COMFORT EZ PEN NEEDLE 12MM | COMFORT TOUCH PEN NDL 31G 8MM |
| 29G157 | |
| COMFORT EZ PEN NEEDLES 4MM | COMFORT TOUCH PEN NDL 32G 4MM |
| 32G SINGLE USE, MICRO 157 | |
| COMFORT EZ PEN NEEDLES 4MM | COMFORT TOUCH PEN NDL 32G 5MM |
| 33G157 | |
| COMFORT EZ PEN NEEDLES 5MM | COMFORT TOUCH PEN NDL 32G 6MM |
| 31G MINI | |
| COMFORT EZ PEN NEEDLES 5MM | COMFORT TOUCH PEN NDL 32G 8MM |
| 32G SINGLE USE,MINI,HRI157 | |
| COMFORT EZ PEN NEEDLES 5MM | COMFORT TOUCH PEN NDL 33G 4MM |
| 33G157 | |
| COMFORT EZ PEN NEEDLES 6MM | COMFORT TOUCH PEN NDL 33G 6MM |
| 31G157 | |
| COMFORT EZ PEN NEEDLES 6MM | COMFORT TOUCH PEN NDL |
| 32G157 | 33GX5MM157 |
| | COPIKTRA99 |
| | |

| COSENTYX (2 SYRINGES)292 | DROPLET 0.5 ML 30GX12.5MM(1/2)157 |
|--|-------------------------------------|
| COSENTYX INTRAVENOUS290 | DROPLET INS 0.3 ML 29GX12.5MM157 |
| COSENTYX PEN (2 PENS)292 | DROPLET INS 0.3 ML 30G 8MM(1/2)157 |
| COSENTYX SUBCUTANEOUS | DROPLET INS 0.3 ML 30GX12.5MM157 |
| SYRINGE 75 MG/0.5 ML292 | DROPLET INS 0.3 ML 31G 6MM(1/2)157 |
| COSENTYX UNOREADY PEN292 | DROPLET INS 0.3 ML 31G 8MM(1/2)157 |
| COTELLIC74 | DROPLET INS 0.5 ML 29G 12.7MM157 |
| CURAD GAUZE PADS 2" X 2" 157 | DROPLET INS 0.5 ML 30G 12.7MM157 |
| CURITY ALCOHOL PREPS 2 | DROPLET INS 0.5 ML 30GX6MM(1/2) 157 |
| PLY,MEDIUM157 | DROPLET INS 0.5 ML 30GX8MM(1/2) 157 |
| CURITY GAUZE SPONGES (12 PLY)- | DROPLET INS 0.5 ML 31GX6MM(1/2) 157 |
| 200/BAG157 | DROPLET INS 0.5 ML 31GX8MM(1/2) 157 |
| CURITY GUAZE PADS 1'S(12 PLY) 157 | DROPLET INS SYR 0.3 ML 30GX6MM. 157 |
| CYLTEZO(CF)17 | DROPLET INS SYR 0.3 ML 30GX8MM. 157 |
| CYLTEZO(CF) PEN17 | DROPLET INS SYR 0.3 ML 31GX6MM. 157 |
| CYLTEZO(CF) PEN CROHN'S-UC-HS17 | DROPLET INS SYR 0.3 ML 31GX8MM. 157 |
| CYLTEZO(CF) PEN PSORIASIS-UV17 | DROPLET INS SYR 0.5 ML 30G 8MM 157 |
| dalfampridine81 | DROPLET INS SYR 0.5 ML 31G 6MM 157 |
| DANYELZA213 | DROPLET INS SYR 0.5 ML 31G 8MM 157 |
| DANZITEN216 | DROPLET INS SYR 1 ML 29GX12.5MM |
| dasatinib oral tablet 100 mg, 140 mg, 20 | |
| mg, 50 mg, 70 mg, 80 mg83 | DROPLET INS SYR 1 ML 30GX12.5MM |
| DATROWAY84 | |
| DAURISMO ORAL TABLET 100 MG, 25 | DROPLET INS SYR 1 ML 30GX6MM 157 |
| MG138 | DROPLET INS SYR 1 ML 30GX8MM 157 |
| deferasirox oral granules in packet86 | DROPLET INS SYR 1 ML 31G 6MM 157 |
| deferasirox oral tablet86 | DROPLET INS SYR 1 ML 31GX6MM 157 |
| DERMACEA 2"X2" GAUZE 12 PLY, | DROPLET INS SYR 1 ML 31GX8MM 157 |
| USP TYPE VII | DROPLET MICRON 34G X 9/64"157 |
| DERMACEA GAUZE 2"X2" SPONGE 8 | DROPLET PEN NEEDLE 29G 10MM 157 |
| PLY157 | DROPLET PEN NEEDLE 29G 12MM 157 |
| DERMACEA NON-WOVEN 2"X2" | DROPLET PEN NEEDLE 30G 8MM 157 |
| SPNGE | DROPLET PEN NEEDLE 31G 5MM 157 |
| dermacinrx lidocan 5% patch outer192 | DROPLET PEN NEEDLE 31G 6MM 157 |
| DIACOMIT ORAL CAPSULE 250 MG, | DROPLET PEN NEEDLE 31G 8MM 157 |
| 500 MG312 | DROPLET PEN NEEDLE 32G 4MM 157 |
| DIACOMIT ORAL POWDER IN | DROPLET PEN NEEDLE 32G 5MM 157 |
| PACKET 250 MG, 500 MG312 | DROPLET PEN NEEDLE 32G 6MM 157 |
| diclofenac sodium topical solution in | DROPLET PEN NEEDLE 32G 8MM 157 |
| metered-dose pump90 | DROPSAFE ALCOHOL 70% PREP |
| dimethyl fumarate oral capsule,delayed | PADS157 |
| release(dr/ec) 120 mg, 120 mg (14)- 240 | DROPSAFE INS SYR 0.3 ML 31G 6MM 157 |
| <i>mg (46), 240 mg</i> 91 | DROPSAFE INS SYR 0.3 ML 31G 8MM 157 |
| dronabinol94 | DROPSAFE INS SYR 0.5 ML 31G 6MM 157 |
| DROPLET 0.3 ML 29G 12.7MM(1/2) 157 | DROPSAFE INS SYR 0.5 ML 31G 8MM 157 |
| DROPLET 0.3 ML 30G 12.7MM(1/2) 157 | DROPSAFE INSUL SYR 1 ML 31G |
| DROPLET 0.5 ML 29GX12.5MM(1/2)157 | 6MM157 |
| | |

| DROPSAFE INSUL SYR 1 ML 31G | EASY TOUCH 1 ML SYR 30GX1/2"157 |
|--|-------------------------------------|
| 8MM157 | EASY TOUCH ALCOHOL 70% PADS |
| DROPSAFE INSULN 1 ML 29G 12.5MM | GAMMA-STERILIZED157 |
| | EASY TOUCH FLIPLOK 1 ML 27GX0.5 157 |
| DROPSAFE PEN NEEDLE 31GX1/4" 157 | EASY TOUCH INSULIN 1 ML 29GX1/2 157 |
| DROPSAFE PEN NEEDLE 31GX3/16" 157 | EASY TOUCH INSULIN 1 ML 30GX1/2 157 |
| DROPSAFE PEN NEEDLE 31GX5/16" 157 | EASY TOUCH INSULIN SYR 0.3 ML 157 |
| <i>droxidopa</i> 95 | EASY TOUCH INSULIN SYR 0.5 ML 157 |
| DRUG MART ULTRA COMFORT SYR.157 | EASY TOUCH INSULIN SYR 1 ML 157 |
| DUPIXENT PEN96 | EASY TOUCH INSULIN SYR 1 ML |
| DUPIXENT SYRINGE96 | RETRACTABLE157 |
| EASY CMFT SFTY PEN NDL 31G 5MM157 | EASY TOUCH INSULN 1 ML 29GX1/2" 157 |
| EASY CMFT SFTY PEN NDL 31G 6MM157 | EASY TOUCH INSULN 1 ML 30GX1/2" 157 |
| EASY CMFT SFTY PEN NDL 32G 4MM157 | EASY TOUCH INSULN 1 ML 30GX5/16157 |
| EASY COMFORT 0.3 ML 31G 1/2"157 | EASY TOUCH INSULN 1 ML 31GX5/16157 |
| EASY COMFORT 0.3 ML 31G 5/16"157 | EASY TOUCH LUER LOK INSUL 1 ML157 |
| EASY COMFORT 0.3 ML SYRINGE 157 | EASY TOUCH PEN NEEDLE 29GX1/2" 157 |
| EASY COMFORT 0.5 ML 30GX1/2"157 | EASY TOUCH PEN NEEDLE 30GX5/16 157 |
| EASY COMFORT 0.5 ML 31GX5/16"157 | EASY TOUCH PEN NEEDLE 31GX1/4" 157 |
| EASY COMFORT 0.5 ML 32GX5/16"157 | EASY TOUCH PEN NEEDLE 31GX3/16 157 |
| EASY COMFORT 0.5 ML SYRINGE 157 | EASY TOUCH PEN NEEDLE 31GX5/16 157 |
| EASY COMFORT 1 ML 31GX5/16"157 | EASY TOUCH PEN NEEDLE 32GX1/4" 157 |
| EASY COMFORT 1 ML 32GX5/16"157 | EASY TOUCH PEN NEEDLE 32GX3/16 157 |
| EASY COMFORT ALCOHOL 70% PAD 157 | EASY TOUCH PEN NEEDLE 32GX5/32 157 |
| EASY COMFORT INSULIN 1 ML SYR157 | EASY TOUCH SAF PEN NDL 29G 5MM |
| EASY COMFORT PEN NDL 31GX1/4" 157 | |
| EASY COMFORT PEN NDL 31GX3/16" 157 | EASY TOUCH SAF PEN NDL 29G 8MM |
| EASY COMFORT PEN NDL 31GX5/16" 157 | |
| EASY COMFORT PEN NDL 32GX5/32" 157 | EASY TOUCH SAF PEN NDL 30G 5MM |
| EASY COMFORT PEN NDL 33G 4MM157 | |
| EASY COMFORT PEN NDL 33G 5MM157 | EASY TOUCH SAF PEN NDL 30G 8MM |
| EASY COMFORT PEN NDL 33G 6MM157 | |
| EASY COMFORT SYR 0.5 ML 29G | EASY TOUCH SYR 0.5 ML 28G |
| 8MM | 12.7MM |
| EASY COMFORT SYR 1 ML 29G 8MM.157 | EASY TOUCH SYR 0.5 ML 29G |
| EASY COMFORT SYR 1 ML 30GX1/2". 157 | 12.7MM |
| EASY GLIDE INS 0.3 ML 31GX6MM 157 | EASY TOUCH SYR 1 ML 27G 16MM 157 |
| EASY GLIDE INS 0.5 ML 31GX6MM 157 | EASY TOUCH SYR 1 ML 28G 12.7MM. 157 |
| EASY GLIDE INS 1 ML 31GX6MM 157 | EASY TOUCH SYR 1 ML 29G 12.7MM. 157 |
| EASY GLIDE PEN NEEDLE 4MM 33G157 | EASY TOUCH UNI-SLIP SYR 1 ML157 |
| EASY TOUCH 0.3 ML SYR 30GX1/2"157 | EASYTOUCH SAF PEN NDL 30G 6MM 157 |
| EASY TOUCH 0.5 ML SYR 27GX1/2"157 | ELAHERE 209 |
| EASY TOUCH 0.5 ML SYR 29GX1/2"157 | ELIGARD |
| EASY TOUCH 0.5 ML SYR 30GX1/2"157 EASY TOUCH 0.5 ML SYR 30GX5/16157 | |
| EASY TOUCH 0.5 ML SYR 30GX3/16 157 EASY TOUCH 1 ML SYR 27GX1/2"157 | ELIGARD (4 MONTH) 185 |
| EASY TOUCH 1 ML SYR 2/GX1/2"157 EASY TOUCH 1 ML SYR 29GX1/2"157 | ELIGARD (6 MONTH) |
| TAST TOUCH LIVIL STR /9UXT//" 13/ | |

| ELREXFIO 44 MG/1.1 ML VIAL INNER, | FIFTY50 PEN 31G X 3/16" NEEDLE |
|---|---|
| SUV, P/F103 | (OTC)157 |
| ELREXFIO SUBCUTANEOUS | <i>fingolimod</i> 129 |
| SOLUTION 40 MG/ML103 | FINTEPLA124 |
| EMBRACE PEN NEEDLE 29G 12MM157 | FOTIVDA332 |
| EMBRACE PEN NEEDLE 30G 5MM157 | FP INSULIN 1 ML SYRINGE157 |
| EMBRACE PEN NEEDLE 30G 8MM157 | FREESTYLE PREC 0.5 ML 30GX5/16 157 |
| EMBRACE PEN NEEDLE 31G 5MM157 | FREESTYLE PREC 0.5 ML 31GX5/16 157 |
| EMBRACE PEN NEEDLE 31G 6MM157 | FREESTYLE PREC 1 ML 30GX5/16" 157 |
| EMBRACE PEN NEEDLE 31G 8MM157 | FREESTYLE PREC 1 ML 31GX5/16" 157 |
| EMBRACE PEN NEEDLE 32G 4MM157 | FRUZAQLA ORAL CAPSULE 1 MG, 5 |
| EMGALITY PEN134 | MG132 |
| EMGALITY SYRINGE | FYARRO301 |
| SUBCUTANEOUS SYRINGE 120 | GAUZE PAD TOPICAL BANDAGE 2 X |
| MG/ML, 300 MG/3 ML (100 MG/ML X | 2 " |
| 3)134 | GAVRETO259 |
| ENBREL117 | gefitinib136 |
| ENBREL MINI117 | GILOTRIF20 |
| ENBREL SURECLICK117 | glatiramer subcutaneous syringe 20 mg/ml, |
| EPCLUSA ORAL PELLETS IN PACKET | <i>40 mg/ml</i> 139 |
| 150-37.5 MG, 200-50 MG303 | glatopa subcutaneous syringe 20 mg/ml, 40 |
| EPCLUSA ORAL TABLET303 | <i>mg/ml</i> |
| EPIDIOLEX64 | glutamine (sickle cell)190 |
| EPKINLY111 | GNP ULT C 0.3 ML 29GX1/2" (1/2) 1/2 |
| EQL INSULIN 0.3 ML SYRINGE | UNIT157 |
| SHORT NEEDLE157 | GNP ULTRA COMFORT 0.5 ML SYR157 |
| EQL INSULIN 0.5 ML SYRINGE | GNP ULTRA COMFORT 1 ML |
| SHORT NEEDLE157 | SYRINGE |
| EQL INSULIN 1 ML SYRINGE SHORT | GNP ULTRA COMFORT 3/10 ML SYR157 |
| NEEDLE157 | GOMEKLI ORAL CAPSULE 1 MG, 2 |
| ERBITUX71 | MG208 |
| ERIVEDGE380 | GOMEKLI ORAL TABLET FOR |
| ERLEADA ORAL TABLET 240 MG, 60 | SUSPENSION208 |
| MG27 | HAEGARDA SUBCUTANEOUS RECON |
| erlotinib oral tablet 100 mg, 150 mg, 25 | SOLN 2,000 UNIT, 3,000 UNIT61 |
| <i>mg</i> 115 | HARVONI ORAL PELLETS IN PACKET |
| everolimus (antineoplastic) oral tablet 10 | 33.75-150 MG, 45-200 MG179 |
| mg, 2.5 mg, 5 mg, 7.5 mg120 | HARVONI ORAL TABLET179 |
| everolimus (antineoplastic) oral tablet for | HEALTHWISE INS 0.3 ML 30GX5/16" 157 |
| suspension121 | HEALTHWISE INS 0.3 ML 31GX5/16" 157 |
| FASENRA49 | HEALTHWISE INS 0.5 ML 30GX5/16" 157 |
| FASENRA PEN49 | HEALTHWISE INS 0.5 ML 31GX5/16" 157 |
| fentanyl citrate buccal lozenge on a handle | HEALTHWISE INS 1 ML 30GX5/16" 157 |
| | HEALTHWISE INS 1 ML 31GX5/16" 157 |
| FIFTY50 INS SYR 1 ML 31GX5/16" | HEALTHWISE PEN NEEDLE 31G 5MM157 |
| SHORT NEEDLE (OTC)157 | HEALTHWISE PEN NEEDLE 31G 8MM157 |
| • | HEALTHWISE PEN NEEDLE 32G 4MM157 |

| HEALTHY ACCENTS PENTIP 4MM | INGREZZA SPRINKLE3 | 72 |
|---|--------------------------------------|-----|
| 32G157 | INLYTA ORAL TABLET 1 MG, 5 MG | 41 |
| HEALTHY ACCENTS PENTIP 5MM | INQOVI | 85 |
| 31G157 | INREBIC1 | 23 |
| HEALTHY ACCENTS PENTIP 6MM | INSULIN SYR 0.3 ML 31GX1/4(1/2)1 | 57 |
| 31G157 | INSULIN SYRIN 0.5 ML 28GX1/2" | |
| HEALTHY ACCENTS PENTIP 8MM | (OTC)1 | 57 |
| 31G157 | INSULIN SYRIN 0.5 ML 29GX1/2" | |
| HEALTHY ACCENTS PENTP 12MM | (OTC)1 | 57 |
| 29G157 | INSULIN SYRIN 0.5 ML 30GX1/2" (RX)1. | |
| HEB INCONTROL ALCOHOL 70% | INSULIN SYRIN 0.5 ML 30GX5/16" | |
| PADS157 | SHORT NEEDLE (OTC)1 | 57 |
| HERCEPTIN HYLECTA348 | INSULIN SYRING 0.5 ML 27G 1/2" | |
| HERZUMA | OUTER1 | |
| HUMIRA PEN11 | INSULIN SYRINGE 0.3 ML1 | 57 |
| HUMIRA PEN CROHNS-UC-HS START. 11 | INSULIN SYRINGE 0.3 ML 31GX1/41 | 57 |
| HUMIRA PEN PSOR-UVEITS-ADOL HS 11 | INSULIN SYRINGE 0.5 ML1 | 57 |
| HUMIRA SUBCUTANEOUS SYRINGE | INSULIN SYRINGE 0.5 ML 31GX1/41 | 57 |
| KIT 40 MG/0.8 ML11 | INSULIN SYRINGE 1 ML1 | 57 |
| HUMIRA(CF)11 | INSULIN SYRINGE 1 ML 27G 1/2" | |
| HUMIRA(CF) PEDI CROHNS STARTER 11 | INNER1 | 57 |
| HUMIRA(CF) PEN11 | INSULIN SYRINGE 1 ML 27G 16MM1 | 57 |
| HUMIRA(CF) PEN CROHNS-UC-HS 11 | INSULIN SYRINGE 1 ML 28GX1/2" | |
| HUMIRA(CF) PEN PEDIATRIC UC 11 | (OTC)1 | 57 |
| HUMIRA(CF) PEN PSOR-UV-ADOL HS. 11 | INSULIN SYRINGE 1 ML 30GX1/2" | |
| IBRANCE240 | (RX)1 | 57 |
| <i>icatibant</i> 148 | INSULIN SYRINGE 1 ML 30GX5/16" | |
| ICLUSIG257 | SHORT NEEDLE (OTC)1 | |
| IDHIFA106 | INSULIN SYRINGE 1 ML 31GX1/4" 1 | 57 |
| imatinib oral tablet 100 mg, 400 mg 150 | INSULIN SYRINGE-NEEDLE U-100 | |
| IMBRUVICA ORAL CAPSULE 140 MG, | SYRINGE 0.3 ML 29 GAUGE, 1 ML 29 | |
| 70 MG147 | GAUGE X 1/2", 1/2 ML 28 GAUGE 1 | 57 |
| IMBRUVICA ORAL SUSPENSION 147 | INSUPEN 30G ULTRAFIN NEEDLE 1 | - / |
| IMBRUVICA ORAL TABLET147 | INSUPEN 31G ULTRAFIN NEEDLE 1 | |
| IMDELLTRA318 | INSUPEN 32G 6MM PEN NEEDLE 1 | |
| IMJUDO352 | INSUPEN 32G 8MM PEN NEEDLE 1 | |
| IMKELDI151 | INSUPEN PEN NEEDLE 29GX12MM 1 | |
| IMPAVIDO207 | INSUPEN PEN NEEDLE 31GX3/16"1 | |
| INCONTROL PEN NEEDLE 12MM 29G 157 | INSUPEN PEN NEEDLE 32GX4MM 1 | |
| INCONTROL PEN NEEDLE 4MM 32G 157 | INSUPEN PEN NEEDLE 33GX4MM 1 | |
| INCONTROL PEN NEEDLE 5MM 31G157 | ITOVEBI ORAL TABLET 3 MG, 9 MG 1 | |
| INCONTROL PEN NEEDLE 6MM 31G157 | IV ANTISEPTIC WIPES1 | |
| INCONTROL PEN NEEDLE 8MM 31G157 | IWILFIN1 | |
| INCRELEX | JAKAFI2 | |
| <i>infliximab</i> 154 | javygtor oral tablet,soluble2 | 89 |
| INGREZZA | JAYPIRCA ORAL TABLET 100 MG, 50 | |
| INGREZZA INITIATION PK(TARDIV)372 | MG2 | 55 |

| JEMPERLI | LITETOUCH SYR 0.5 ML 29GX1/2" 157 |
|--|--------------------------------------|
| KALYDECO172 | LITETOUCH SYR 0.5 ML 30GX5/16" 157 |
| KENDALL ALCOHOL 70% PREP PAD. 157 | LITETOUCH SYRIN 1 ML 28GX1/2" 157 |
| KERENDIA128 | LITETOUCH SYRIN 1 ML 29GX1/2" 157 |
| KESIMPTA PEN229 | LITETOUCH SYRIN 1 ML 30GX5/16" 157 |
| KEYTRUDA248 | LIVTENCITY200 |
| KIMMTRAK320 | LONSURF ORAL TABLET 15-6.14 MG, |
| KINERET25 | 20-8.19 MG |
| KISQALI FEMARA CO-PACK ORAL | LOQTORZI342 |
| TABLET 200 MG/DAY(200 MG X 1)-2.5 | LORBRENA ORAL TABLET 100 MG, |
| MG, 400 MG/DAY(200 MG X 2)-2.5 MG, | 25 MG |
| 600 MG/DAY(200 MG X 3)-2.5 MG 271 | LUMAKRAS ORAL TABLET 120 MG, |
| KISQALI ORAL TABLET 200 MG/DAY | 240 MG, 320 MG311 |
| (200 MG X 1), 400 MG/DAY (200 MG X | LUNSUMIO211 |
| 2), 600 MG/DAY (200 MG X 3)270 | LUPRON DEPOT 186 |
| KOSELUGO ORAL CAPSULE 10 MG, | LUPRON DEPOT (3 MONTH)186 |
| 25 MG298 | LUPRON DEPOT (4 MONTH)186 |
| KRAZATI10 | LUPRON DEPOT (6 MONTH)186 |
| KYNMOBI SUBLINGUAL FILM 10 MG, | LUPRON DEPOT-PED (3 MONTH) 188 |
| 10-15-20-25-30 MG, 15 MG, 20 MG, 25 | LUPRON DEPOT-PED |
| MG, 30 MG29 | INTRAMUSCULAR SYRINGE KIT188 |
| lanreotide subcutaneous syringe 120 | LYBALVI230 |
| <i>mg/0.5 ml</i> 175 | LYNPARZA231 |
| <i>lapatinib</i> 176 | LYTGOBI ORAL TABLET 12 MG/DAY |
| LAZCLUZE ORAL TABLET 240 MG, 80 | (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 |
| MG178 | MG/DAY (4 MG X 5)133 |
| lenalidomide180 | MAGELLAN INSUL SYRINGE 0.3 ML157 |
| LENVIMA181 | MAGELLAN INSUL SYRINGE 0.5 ML157 |
| <i>leuprolide (3 month)</i> 184 | MAGELLAN INSULIN SYR 0.3 ML 157 |
| leuprolide subcutaneous kit183 | MAGELLAN INSULIN SYR 0.5 ML 157 |
| lidocaine topical adhesive patch,medicated | MAGELLAN INSULIN SYRINGE 1 ML 157 |
| 5 %192 | MARGENZA199 |
| lidocaine topical ointment191 | MAVENCLAD (10 TABLET PACK)72 |
| lidocaine-prilocaine topical cream 193 | MAVENCLAD (4 TABLET PACK)72 |
| lidocan iii192 | MAVENCLAD (5 TABLET PACK)72 |
| LISCO SPONGES 100/BAG157 | MAVENCLAD (6 TABLET PACK)72 |
| LITE TOUCH 31GX1/4" PEN NEEDLE 157 | MAVENCLAD (7 TABLET PACK)72 |
| LITE TOUCH INSULIN 0.5 ML SYR 157 | MAVENCLAD (8 TABLET PACK)72 |
| LITE TOUCH INSULIN 1 ML SYR157 | MAVENCLAD (9 TABLET PACK)72 |
| LITE TOUCH INSULIN SYR 1 ML157 | MAXICOMFORT II PEN NDL |
| LITE TOUCH PEN NEEDLE 29G157 | 31GX6MM |
| LITE TOUCH PEN NEEDLE 31G157 | MAXICOMFORT INS 0.5 ML 27GX1/2" 157 |
| LITETOUCH INS 0.3 ML 29GX1/2"157 | MAXI-COMFORT INS 0.5 ML 28G157 |
| LITETOUCH INS 0.3 ML 30GX5/16"157 | MAXICOMFORT INS 1 ML 27GX1/2" 157 |
| LITETOUCH INS 0.3 ML 31GX5/16"157 | MAXI-COMFORT INS 1 ML 28GX1/2"157 |
| LITETOUCH INS 0.5 ML 31GX5/16"157 | |
| LITETOLICH SVR 0.5 ML 28GX1/2" 157 | |

| MAXICOMFORT PEN NDL 29G X 5MM | MONOJECT INSULIN SYR 0.3 ML | |
|--|------------------------------------|-------|
| | (OTC) | .157 |
| MAXICOMFORT PEN NDL 29G X 8MM | MONOJECT INSULIN SYR 0.5 ML | . 157 |
| | MONOJECT INSULIN SYR 0.5 ML | |
| MAYZENT ORAL TABLET 0.25 MG, 1 | (OTC) | .157 |
| MG, 2 MG300 | MONOJECT INSULIN SYR 1 ML 3'S | |
| MAYZENT STARTER(FOR 1MG | (OTC) | .157 |
| MAINT)300 | MONOJECT INSULIN SYR U-100 | |
| MAYZENT STARTER(FOR 2MG | MONOJECT SYRINGE 0.3 ML | .157 |
| MAINT)300 | MONOJECT SYRINGE 0.5 ML | .157 |
| MEKINIST ORAL RECON SOLN 344 | MONOJECT SYRINGE 1 ML | .157 |
| MEKINIST ORAL TABLET 0.5 MG, 2 | morphine concentrate oral solution | 146 |
| MG345 | MOUNJARO | |
| MEKTOVI56 | MVASI | 53 |
| MICRODOT PEN NEEDLE 31GX6MM157 | NANO 2 GEN PEN NEEDLE 32G 4MM. | 157 |
| MICRODOT PEN NEEDLE 32GX4MM157 | NATPARA | |
| MICRODOT PEN NEEDLE 33GX4MM157 | NERLYNX | 214 |
| MICRODOT READYGARD NDL 31G | NEULASTA ONPRO | .245 |
| 5MM OUTER157 | NIKTIMVO | 40 |
| mifepristone oral tablet 300 mg 206 | NINLARO | 174 |
| MINI PEN NEEDLE 32G 4MM157 | nitisinone | 222 |
| MINI PEN NEEDLE 32G 5MM157 | NIVESTYM | |
| MINI PEN NEEDLE 32G 6MM157 | NORDITROPIN FLEXPRO | 305 |
| MINI PEN NEEDLE 32G 8MM157 | NOVOFINE 30 | |
| MINI PEN NEEDLE 33G 4MM157 | NOVOFINE 32G NEEDLES | . 157 |
| MINI PEN NEEDLE 33G 5MM157 | NOVOFINE PLUS PEN NDL 32GX1/6". | .157 |
| MINI PEN NEEDLE 33G 6MM157 | NOVOTWIST NEEDLE 32G 5MM | .157 |
| MINI ULTRA-THIN II PEN NDL 31G | NUBEQA | 82 |
| STERILE | NUCALA SUBCUTANEOUS AUTO- | |
| MIPLYFFA | INJECTOR | 203 |
| modafinil oral tablet 100 mg, 200 mg 212 | NUCALA SUBCUTANEOUS RECON | |
| MONOJECT 0.5 ML SYRN 28GX1/2"157 | SOLN | |
| MONOJECT 1 ML SYRN 27X1/2"157 | NUCALA SUBCUTANEOUS SYRINGE | |
| MONOJECT 1 ML SYRN 28GX1/2" | 100 MG/ML, 40 MG/0.4 ML | |
| (OTC)157 | NUPLAZID | |
| MONOJECT INSUL SYR U100 (OTC)157 | NURTEC ODT | |
| MONOJECT INSUL SYR U100 | NYVEPRIA | |
| .5ML,29GX1/2" (OTC)157 | OCREVUS | |
| MONOJECT INSUL SYR U100 0.5 ML | OCREVUS ZUNOVO | |
| CONVERTS TO 29G (OTC)157 | ODOMZO | |
| MONOJECT INSUL SYR U100 1 ML 157 | OFEV | |
| MONOJECT INSUL SYR U100 1 ML 3'S, | OGIVRI | |
| 29GX1/2" (OTC) | OGSIVEO ORAL TABLET 100 MG, 150 | |
| MONOJECT INSUL SYR U100 1 ML | MG, 50 MG | 221 |
| W/O NEEDLE (OTC) | OJEMDA ORAL SUSPENSION FOR | |
| MONOJECT INSULIN SYR 0.3 ML 157 | RECONSTITUTION | |
| | OJEMDA ORAL TABLET | .343 |

| OJJAARA210 | PENTIPS PEN NEEDLE 32GX5/32" |
|----------------------------------|------------------------------------|
| ONAPGO28 | 4MM157 |
| ONTRUZANT347 | PIP PEN NEEDLE 31G X 5MM 157 |
| ONUREG42 | PIP PEN NEEDLE 32G X 4MM 157 |
| OPDIVO223 | |
| OPDIVO QVANTIG224 | (200 MG X 1), 250 MG/DAY (200 MG |
| OPDUALAG225 | |
| OPSUMIT198 | 2)22 |
| ORENCIA4 | pirfenidone oral capsule254 |
| ORENCIA (WITH MALTOSE)2 | |
| ORENCIA CLICKJECT4 | 801 mg254 |
| ORFADIN ORAL SUSPENSION222 | PLEGRIDY SUBCUTANEOUS PEN |
| ORGOVYX264 | |
| ORILISSA ORAL TABLET 150 MG, 200 | MCG/0.5 ML- 94 MCG/0.5 ML 169 |
| MG102 | PLEGRIDY SUBCUTANEOUS |
| ORKAMBI ORAL TABLET197 | SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 |
| ORSERDU ORAL TABLET 345 MG, 86 | ML- 94 MCG/0.5 ML169 |
| MG101 | POMALYST256 |
| OTEZLA30 | |
| OTEZLA STARTER30 | |
| oxandrolone238 | PREVENT PEN NEEDLE 31GX1/4"157 |
| OZEMPIC141 | PREVENT PEN NEEDLE 31GX5/16"157 |
| pazopanib243 | |
| PC UNIFINE PENTIPS 8MM NEEDLE | PRO COMFORT 0.5 ML 30GX1/2"157 |
| SHORT | |
| PEGASYS246 | |
| PEMAZYRE249 | |
| PEN NEEDLE 30G 5MM OUTER157 | |
| PEN NEEDLE 30G 8MM INNER157 | |
| PEN NEEDLE 30G X 5/16"157 | |
| PEN NEEDLE, DIABETIC NEEDLE 29 | |
| GAUGE X 1/2"157 | |
| PEN NEEDLES 12MM 29G | PRO COMFORT PEN NDL 4MM 32G 157 |
| 29GX12MM,STRL | PRO COMFORT PEN NDL 5MM 32G 157 |
| PEN NEEDLES 4MM 32G157 | |
| PEN NEEDLES 6MM 31G 31GX6MM, | PRODIGY SYRNG 0.5 ML 31GX5/16" 157 |
| STRL | |
| PEN NEEDLES 8MM 31G | PROMACTA ORAL POWDER IN |
| 31GX8MM,STRL,SHORT (OTC)157 | |
| penicillamine oral tablet250 | · |
| PENTIPS PEN NEEDLE 29G 1/2"157 | , , |
| PENTIPS PEN NEEDLE 31G 1/4"157 | |
| PENTIPS PEN NEEDLE 31GX3/16" | PURE CMFT SFTY PEN NDL 31G 6MM157 |
| MINI, 5MM | |
| PENTIPS PEN NEEDLE 31GX5/16" | PURE COMFORT ALCOHOL 70% |
| SHORT, 8MM | |
| PENTIPS PEN NEEDLE 32G 1/4"157 | PURE COMFORT PEN NDL 32G 4MM157 |

| PURE COMFORT PEN NDL 32G 5MM157 | SAFESNAP INS SYR UNITS-100 0.3 ML |
|--------------------------------------|--|
| PURE COMFORT PEN NDL 32G 6MM157 | 30GX5/16",10X10157 |
| PURE COMFORT PEN NDL 32G 8MM157 | SAFESNAP INS SYR UNITS-100 0.5 ML |
| pyrimethamine260 | 29GX1/2",10X10157 |
| QINLOCK279 | SAFESNAP INS SYR UNITS-100 0.5 ML |
| quinine sulfate261 | 30GX5/16",10X10157 |
| QULIPTA37 | SAFESNAP INS SYR UNITS-100 1 ML |
| RALDESY351 | 28GX1/2",10X10157 |
| RAYA SURE PEN NEEDLE 29G 12MM.157 | SAFESNAP INS SYR UNITS-100 1 ML |
| RAYA SURE PEN NEEDLE 31G 4MM157 | 29GX1/2",10X10157 |
| RAYA SURE PEN NEEDLE 31G 5MM157 | SAFETY PEN NEEDLE 31G 4MM157 |
| RAYA SURE PEN NEEDLE 31G 6MM157 | SAFETY PEN NEEDLE 5MM X 31G 157 |
| RELION INS SYR 0.3 ML 31GX6MM 157 | SAFETY SYRINGE 0.5 ML 30G 1/2" 157 |
| RELION INS SYR 0.5 ML 31GX6MM 157 | sapropterin oral tablet, soluble |
| RELION INS SYR 1 ML 31GX15/64"157 | SCEMBLIX ORAL TABLET 100 MG, 20 |
| RELI-ON INSULIN 0.5 ML SYR 157 | MG, 40 MG |
| RELI-ON INSULIN 1 ML SYR 157 | SECURESAFE PEN NDL 30GX5/16" |
| RELION MINI PEN 31G X 1/4" NDL 157 | OUTER157 |
| RETACRIT INJECTION SOLUTION | SECURESAFE SYR 0.5 ML 29G 1/2" |
| 10,000 UNIT/ML, 2,000 UNIT/ML, | OUTER |
| 20,000 UNIT/2 ML, 20,000 UNIT/ML, | SECURESAFE SYRNG 1 ML 29G 1/2" |
| 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 | OUTER |
| UNIT/ML | SELARSDI |
| RETEVMO ORAL CAPSULE 40 MG, 80 | SEROSTIM SUBCUTANEOUS RECON |
| MG297 | SOLN 4 MG, 5 MG, 6 MG |
| RETEVMO ORAL TABLET 120 MG, 160 | SIGNIFOR 242 |
| MG, 40 MG, 80 MG297 | sildenafil (pulm.hypertension) oral tablet 299 |
| REVUFORJ ORAL TABLET 110 MG, | SIRTURO44 |
| 160 MG, 25 MG269 | SKY SAFETY PEN NEEDLE 30G 5MM.157 |
| REZLIDHIA232 | SKY SAFETY PEN NEEDLE 30G 8MM.157 |
| REZUROCK46 | SKYRIZI |
| RIABNI | SM ULT CFT 0.3 ML 31GX5/16(1/2)157 |
| RINVOQ358 | sodium oxybate302 |
| RINVOQ LQ | SOMATULINE DEPOT |
| RITUXAN HYCELA282 | SUBCUTANEOUS SYRINGE 60 MG/0.2 |
| ROMVIMZA379 | ML, 90 MG/0.3 ML175 |
| ROZLYTREK ORAL CAPSULE 100 MG, | SOMAVERT247 |
| 200 MG108 | sorafenib309 |
| ROZLYTREK ORAL PELLETS IN | SPRAVATO116 |
| PACKET109 | STELARA |
| RUBRACA287 | STERILE PADS 2" X 2" |
| RUXIENCE285 | STIVARGA263 |
| RYBELSUS141 | STRENSIQ34 |
| RYBREVANT24 | sunitinib malate313 |
| RYDAPT205 | SURE CMFT SFTY PEN NDL 31G 6MM157 |
| RYTELO | SURE CMFT SFTY PEN NDL 32G 4MM157 |
| | SURE COMFORT 0.5 ML SYRINGE 157 |
| | |

| SURE COMFORT 1 ML SYRINGE 157 | TECHLITE INS SYR 1 ML 29GX12MM.15 |
|---------------------------------------|--|
| SURE COMFORT 3/10 ML SYRINGE 157 | TECHLITE INS SYR 1 ML 30GX12MM.15 |
| SURE COMFORT 3/10 ML SYRINGE | TECHLITE INS SYR 1 ML 30GX8MM15 |
| INSULIN SYRINGE157 | TECHLITE INS SYR 1 ML 31GX6MM15 |
| SURE COMFORT 30G PEN NEEDLE157 | TECHLITE INS SYR 1 ML 31GX8MM15 |
| SURE COMFORT ALCOHOL PREP | TECHLITE PEN NEEDLE 29GX1/2" 15 |
| PADS157 | TECHLITE PEN NEEDLE 29GX3/8" 15 |
| SURE COMFORT INS 0.3 ML 31GX1/4.157 | TECHLITE PEN NEEDLE 31GX1/4" 15 |
| SURE COMFORT INS 0.5 ML 31GX1/4.157 | TECHLITE PEN NEEDLE 31GX3/16" 15 |
| SURE COMFORT INS 1 ML 31GX1/4"157 | TECHLITE PEN NEEDLE 31GX5/16" 15 |
| SURE COMFORT PEN NDL 29GX1/2" | TECHLITE PEN NEEDLE 32GX1/4" 15 |
| 12.7MM157 | TECHLITE PEN NEEDLE 32GX5/16" 15 |
| SURE COMFORT PEN NDL 31G 5MM157 | TECHLITE PEN NEEDLE 32GX5/32" 15 |
| SURE COMFORT PEN NDL 31G 8MM157 | TECHLITE PLUS PEN NDL 32G 4MM15 |
| SURE COMFORT PEN NDL 32G 4MM157 | TECVAYLI32 |
| SURE COMFORT PEN NDL 32G 6MM157 | TEPMETKO32 |
| SURE-FINE PEN NEEDLES 12.7MM 157 | teriparatide subcutaneous pen injector 20 |
| SURE-FINE PEN NEEDLES 5MM 157 | mcg/dose (620mcg/2.48ml)324 |
| SURE-FINE PEN NEEDLES 8MM 157 | TERUMO INS SYRINGE U100-1 ML 15' |
| SURE-JECT INSU SYR U100 0.3 ML 157 | TERUMO INS SYRINGE U100-1/2 ML 15 |
| SURE-JECT INSU SYR U100 0.5 ML 157 | TERUMO INS SYRINGE U100-1/3 ML 15 |
| SURE-JECT INSU SYR U100 1 ML 157 | TERUMO INS SYRNG U100-1/2 ML 15 |
| SURE-JECT INSUL SYR U100 1 ML 157 | testosterone cypionate32 |
| SURE-JECT INSULIN SYRINGE 1 ML 157 | testosterone enanthate32 |
| SURE-PREP ALCOHOL PREP PADS 157 | testosterone transdermal gel in metered- |
| SYMPAZAN73 | dose pump 12.5 mg/ 1.25 gram (1 %), |
| SYNRIBO233 | 20.25 mg/1.25 gram (1.62 %)32. |
| TABRECTA66 | testosterone transdermal gel in packet 1 % |
| tadalafil oral tablet 2.5 mg, 5 mg315 | (25 mg/2.5gram), 1 % (50 mg/5 gram)32 |
| TAFINLAR ORAL CAPSULE78 | tetrabenazine32 |
| TAFINLAR ORAL TABLET FOR | TEVIMBRA330 |
| SUSPENSION79 | THALOMID329 |
| TAGRISSO237 | THINPRO INS SYRIN U100-0.3 ML 15 |
| TALVEY317 | THINPRO INS SYRIN U100-0.5 ML 15 |
| TALZENNA316 | THINPRO INS SYRIN U100-1 ML 15 |
| TASIGNA ORAL CAPSULE 150 MG, | TIBSOVO173 |
| 200 MG, 50 MG215 | TIVDAK33 |
| TAVNEOS38 | TOPCARE CLICKFINE 31G X 1/4"15 |
| TAZVERIK319 | TOPCARE CLICKFINE 31G X 5/16"15 |
| TECHLITE 0.3 ML 29GX12MM (1/2) 157 | TOPCARE ULTRA COMFORT |
| TECHLITE 0.3 ML 30GX8MM (1/2) 157 | SYRINGE15 |
| TECHLITE 0.3 ML 31GX6MM (1/2) 157 | torpenz oral tablet 10 mg, 2.5 mg, 5 mg, |
| TECHLITE 0.3 ML 31GX8MM (1/2) 157 | 7.5 mg |
| TECHLITE 0.5 ML 30GX12MM (1/2) 157 | TRAZIMERA350 |
| TECHLITE 0.5 ML 30GX8MM (1/2) 157 | TRELSTAR INTRAMUSCULAR |
| TECHLITE 0.5 ML 31GX6MM (1/2) 157 | SUSPENSION FOR RECONSTITUTION 35. |
| TECHLITE 0.5 ML 31GX8MM (1/2) 157 | TREMFYA14 |

| TREMFYA PEN SUBCUTANEOUS PEN | TRUEPLUS SYR 0.5 ML 30GX5/16" | .157 |
|-------------------------------------|----------------------------------|-------|
| INJECTOR 200 MG/2 ML144 | TRUEPLUS SYR 0.5 ML 31GX5/16" | .157 |
| tretinoin topical cream341 | TRUEPLUS SYR 1 ML 28GX1/2" | .157 |
| trientine oral capsule 250 mg353 | TRUEPLUS SYR 1 ML 29GX1/2" | .157 |
| TRUE CMFRT PRO 0.5 ML 30G 5/16"157 | TRUEPLUS SYR 1 ML 30GX5/16" | .157 |
| TRUE CMFRT PRO 0.5 ML 31G 5/16"157 | TRUEPLUS SYR 1 ML 31GX5/16" | .157 |
| TRUE CMFRT PRO 0.5 ML 32G 5/16"157 | TRULICITY | 140 |
| TRUE CMFT SFTY PEN NDL 31G 5MM157 | TRUQAP | |
| TRUE CMFT SFTY PEN NDL 31G 6MM157 | TRUXIMA | |
| TRUE CMFT SFTY PEN NDL 32G 4MM157 | TUKYSA ORAL TABLET 150 MG, 50 | |
| TRUE COMFORT 0.5 ML 30G 1/2"157 | MG | .356 |
| TRUE COMFORT 0.5 ML 30G 5/16"157 | TURALIO | .252 |
| TRUE COMFORT 0.5 ML 31G 5/16"157 | TYENNE | |
| TRUE COMFORT 0.5 ML 31GX5/16"157 | TYENNE AUTOINJECTOR | 337 |
| TRUE COMFORT 1 ML 31GX5/16"157 | TYMLOS | 1 |
| TRUE COMFORT ALCOHOL 70% | UBRELVY | |
| PADS157 | ULTICAR INS 0.3 ML 31GX1/4(1/2) | .157 |
| TRUE COMFORT PEN NDL 31G 8MM157 | ULTICARE INS 1 ML 31GX1/4" | |
| TRUE COMFORT PEN NDL 31GX5MM 157 | ULTICARE INS SYR 0.3 ML 30G 8MM. | 157 |
| TRUE COMFORT PEN NDL 31GX6MM 157 | ULTICARE INS SYR 0.3 ML 31G 6MM. | 157 |
| TRUE COMFORT PEN NDL 32G 5MM157 | ULTICARE INS SYR 0.3 ML 31G 8MM. | 157 |
| TRUE COMFORT PEN NDL 32G 6MM157 | ULTICARE INS SYR 0.5 ML 31G 6MM. | 157 |
| TRUE COMFORT PEN NDL 32GX4MM 157 | ULTICARE INS SYR 0.5 ML 31G 8MM | |
| TRUE COMFORT PEN NDL 33G 4MM157 | (OTC) | .157 |
| TRUE COMFORT PEN NDL 33G 5MM157 | ULTICARE INS SYR 1 ML 30GX1/2" | .157 |
| TRUE COMFORT PEN NDL 33G 6MM157 | ULTICARE PEN NEEDLE 31GX3/16" | .157 |
| TRUE COMFORT PRO 1 ML 30G 1/2"157 | ULTICARE PEN NEEDLE 6MM 31G | 157 |
| TRUE COMFORT PRO 1 ML 30G 5/16".157 | ULTICARE PEN NEEDLE 8MM 31G | 157 |
| TRUE COMFORT PRO 1 ML 31G 5/16".157 | ULTICARE PEN NEEDLES 12MM 29G | . 157 |
| TRUE COMFORT PRO 1 ML 32G 5/16".157 | ULTICARE PEN NEEDLES 4MM 32G | |
| TRUE COMFORT PRO ALCOHOL | MICRO, 32GX4MM | |
| PADS157 | ULTICARE PEN NEEDLES 6MM 32G | . 157 |
| TRUE COMFORT SFTY 1 ML 30G 1/2".157 | ULTICARE SAFE PEN NDL 30G 8MM. | .157 |
| TRUE COMFRT PRO 0.5 ML 30G 1/2"157 | ULTICARE SAFE PEN NDL 5MM 30G. | |
| TRUE COMFRT SFTY 1 ML 30G 5/16"157 | ULTICARE SYR 0.3 ML 29G 12.7MM | .157 |
| TRUE COMFRT SFTY 1 ML 31G 5/16"157 | ULTICARE SYR 0.3 ML 30GX1/2" | . 157 |
| TRUE COMFRT SFTY 1 ML 32G 5/16"157 | ULTICARE SYR 0.3 ML 31GX5/16" | |
| TRUEPLUS PEN NEEDLE 29GX1/2" 157 | SHORT NDL | |
| TRUEPLUS PEN NEEDLE 31G X 1/4" 157 | ULTICARE SYR 0.5 ML 30GX1/2" | . 157 |
| TRUEPLUS PEN NEEDLE 31GX3/16" 157 | ULTICARE SYR 0.5 ML 31GX5/16" | |
| TRUEPLUS PEN NEEDLE 31GX5/16" 157 | SHORT NDL | |
| TRUEPLUS PEN NEEDLE 32GX5/32" 157 | ULTICARE SYR 1 ML 31GX5/16" | |
| TRUEPLUS SYR 0.3 ML 29GX1/2"157 | ULTIGUARD SAFE 1 ML 30G 12.7MM | . 157 |
| TRUEPLUS SYR 0.3 ML 30GX5/16"157 | ULTIGUARD SAFE0.3 ML 30G 12.7MM | |
| TRUEPLUS SYR 0.3 ML 31GX5/16"157 | | 157 |
| TRUEPLUS SYR 0.5 ML 28GX1/2"157 | | |
| TRUEPLUS SVR 0.5 ML 29GX1/2" 157 | | |

| III TIGILADD CAFEO CAR ANG 12 TABLE | THE ACADE DIG 1 MI 200M / OH |
|-------------------------------------|-------------------------------------|
| ULTIGUARD SAFE0.5 ML 30G 12.7MM | ULTRACARE INS 1 ML 30GX1/2" 157 |
| | ULTRACARE INS 1 ML 31G X 5/16" 157 |
| ULTIGUARD SAFEPACK 1 ML 31G | ULTRACARE PEN NEEDLE 31GX1/4"157 |
| 8MM157 | ULTRACARE PEN NEEDLE 31GX3/16"157 |
| ULTIGUARD SAFEPACK 29G 12.7MM 157 | ULTRACARE PEN NEEDLE 31GX5/16"157 |
| ULTIGUARD SAFEPACK 31G 5MM 157 | ULTRACARE PEN NEEDLE 32GX1/4"157 |
| | |
| ULTIGUARD SAFEPACK 31G 6MM 157 | ULTRACARE PEN NEEDLE 32GX3/16" 157 |
| ULTIGUARD SAFEPACK 31G 8MM 157 | ULTRACARE PEN NEEDLE 32GX5/32"157 |
| ULTIGUARD SAFEPACK 32G 4MM 157 | ULTRACARE PEN NEEDLE 33GX5/32"157 |
| ULTIGUARD SAFEPACK 32G 6MM 157 | ULTRA-FINE 0.3 ML 30G 12.7MM157 |
| ULTIGUARD SAFEPK 0.3 ML 31G | ULTRA-FINE 0.3 ML 31G 6MM (1/2)157 |
| 8MM157 | ULTRA-FINE 0.3 ML 31G 8MM (1/2)157 |
| ULTIGUARD SAFEPK 0.5 ML 31G | ULTRA-FINE 0.5 ML 30G 12.7MM157 |
| | |
| 8MM | ULTRA-FINE INS SYR 1 ML 31G 8MM 157 |
| ULTILET ALCOHOL STERL SWAB 157 | ULTRA-FINE PEN NDL 29G 12.7MM157 |
| ULTILET INSULIN SYRINGE 0.3 ML 157 | ULTRA-FINE PEN NEEDLE 32G 6MM157 |
| ULTILET INSULIN SYRINGE 0.5 ML 157 | ULTRA-FINE SYR 0.5 ML 31G 8MM 157 |
| ULTILET INSULIN SYRINGE 1 ML 157 | ULTRA-FINE SYR 1 ML 30G 12.7MM157 |
| ULTILET PEN NEEDLE157 | ULTRA-THIN II 1 ML 31GX5/16"157 |
| ULTILET PEN NEEDLE 4MM 32G157 | ULTRA-THIN II INS 0.3 ML 30G157 |
| | |
| ULTRA COMFORT 0.3 ML SYRINGE 157 | ULTRA-THIN II INS 0.3 ML 31G157 |
| ULTRA COMFORT 0.5 ML 28GX1/2" | ULTRA-THIN II INS 0.5 ML 29G157 |
| CONVERTS TO 29G157 | ULTRA-THIN II INS 0.5 ML 30G157 |
| ULTRA COMFORT 0.5 ML 29GX1/2" 157 | ULTRA-THIN II INS 0.5 ML 31G157 |
| ULTRA COMFORT 0.5 ML SYRINGE 157 | ULTRA-THIN II INS SYR 1 ML 29G 157 |
| ULTRA COMFORT 1 ML 31GX5/16"157 | ULTRA-THIN II INS SYR 1 ML 30G 157 |
| ULTRA COMFORT 1 ML SYRINGE 157 | ULTRA-THIN II PEN NDL 29GX1/2" 157 |
| | |
| ULTRA FLO 0.3 ML 30G 1/2" (1/2)157 | ULTRA-THIN II PEN NDL 31GX5/16157 |
| ULTRA FLO 0.3 ML 30G 5/16"(1/2)157 | UNIFINE OTC PEN NEEDLE 31G 5MM 157 |
| ULTRA FLO 0.3 ML 31G 5/16"(1/2)157 | UNIFINE OTC PEN NEEDLE 32G 4MM 157 |
| ULTRA FLO PEN NEEDLE 31G 5MM 157 | UNIFINE PEN NEEDLE 32G 4MM 157 |
| ULTRA FLO PEN NEEDLE 31G 8MM 157 | UNIFINE PENTIPS 12MM 29G |
| ULTRA FLO PEN NEEDLE 32G 4MM 157 | 29GX12MM, STRL |
| ULTRA FLO PEN NEEDLE 33G 4MM 157 | UNIFINE PENTIPS 31GX3/16" |
| ULTRA FLO PEN NEEDLES 12MM 29G | 31GX5MM,STRL,MINI |
| | |
| | UNIFINE PENTIPS 32GX1/4"157 |
| ULTRA FLO SYR 0.3 ML 29GX1/2" 157 | UNIFINE PENTIPS 32GX5/32" |
| ULTRA FLO SYR 0.3 ML 30G 5/16"157 | 32GX4MM, STRL, NANO157 |
| ULTRA FLO SYR 0.3 ML 31G 5/16"157 | UNIFINE PENTIPS 33GX5/32"157 |
| ULTRA FLO SYR 0.5 ML 29G 1/2"157 | UNIFINE PENTIPS 6MM 31G 157 |
| ULTRA THIN PEN NDL 32G X 4MM157 | UNIFINE PENTIPS MAX 30GX3/16"157 |
| ULTRACARE INS 0.3 ML 30GX5/16" 157 | UNIFINE PENTIPS NEEDLES 29G157 |
| ULTRACARE INS 0.3 ML 31GX5/16" 157 | UNIFINE PENTIPS PLUS 29GX1/2" |
| | |
| ULTRACARE INS 0.5 ML 30GX1/2" 157 | 12MM |
| ULTRACARE INS 0.5 ML 30GX5/16" 157 | UNIFINE PENTIPS PLUS 30GX3/16" 157 |
| ULTRACARE INS 0.5 ML 31GX5/16" 157 | UNIFINE PENTIPS PLUS 31GX1/4" |
| ULTRACARE INS 1 ML 30G X 5/16" 157 | ULTRA SHORT, 6MM157 |
| | |

| UNIFINE PENTIPS PLUS 31GX3/16" | VERIFINE PLUS PEN NDL 32G 4MM- | |
|--------------------------------------|----------------------------------|-------|
| MINI | SHARPS CONTAINER | 157 |
| UNIFINE PENTIPS PLUS 31GX5/16" | VERIFINE SYRING 0.5 ML 29G 1/2" | . 157 |
| SHORT157 | VERIFINE SYRING 1 ML 31G 5/16" | |
| UNIFINE PENTIPS PLUS 32GX5/32" 157 | VERIFINE SYRNG 0.3 ML 31G 5/16" | 157 |
| UNIFINE PENTIPS PLUS 33GX5/32" 157 | VERIFINE SYRNG 0.5 ML 31G 5/16" | 157 |
| UNIFINE PROTECT 30G 5MM 157 | VERQUVO | 377 |
| UNIFINE PROTECT 30G 8MM 157 | VERSALON ALL PURPOSE SPONGE | |
| UNIFINE PROTECT 32G 4MM 157 | 25'S,N-STERILE,3PLY | . 157 |
| UNIFINE SAFECONTROL 30G 5MM157 | VERZENIO | |
| UNIFINE SAFECONTROL 30G 8MM157 | vigabatrin | |
| UNIFINE SAFECONTROL 31G 5MM157 | vigadrone | |
| UNIFINE SAFECONTROL 31G 6MM157 | vigpoder | |
| UNIFINE SAFECONTROL 31G 8MM157 | VITRAKVI ORAL CAPSULE 100 MG, | |
| UNIFINE SAFECONTROL 32G 4MM157 | 25 MG | 177 |
| UNIFINE ULTRA PEN NDL 31G 5MM 157 | VITRAKVI ORAL SOLUTION | |
| UNIFINE ULTRA PEN NDL 31G 6MM 157 | VIZIMPRO | |
| UNIFINE ULTRA PEN NDL 31G 8MM 157 | VONJO | |
| UNIFINE ULTRA PEN NDL 32G 4MM 157 | VORANIGO | |
| UPTRAVI INTRAVENOUS295 | voriconazole oral suspension for | |
| UPTRAVI ORAL TABLET 1,000 MCG, | reconstitution | 382 |
| 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 | VOSEVI | |
| MCG, 400 MCG, 600 MCG, 800 MCG295 | VOWST | 122 |
| UPTRAVI ORAL TABLETS,DOSE | VUMERITY | 92 |
| PACK | VYALEV | 130 |
| VALCHLOR | VYLOY | 386 |
| VANFLYTA262 | WEBCOL ALCOHOL PREPS | |
| VANISHPOINT 0.5 ML 30GX1/2" SY | 20'S,LARGE | 157 |
| OUTER157 | WELIREG | |
| VANISHPOINT INS 1 ML 30GX3/16" 157 | WINREVAIR | . 310 |
| VANISHPOINT U-100 29X1/2 SYR 157 | XALKORI ORAL CAPSULE | 76 |
| VEGZELMA52 | XALKORI ORAL PELLET 150 MG, 20 | |
| VENCLEXTA ORAL TABLET 10 MG, | MG, 50 MG | 77 |
| 100 MG, 50 MG376 | XDEMVY | |
| VENCLEXTA STARTING PACK376 | XELJANZ | 339 |
| VEOZAH126 | XELJANZ XR | 339 |
| VERIFINE INS SYR 1 ML 29G 1/2" 157 | XERMELO | 322 |
| VERIFINE PEN NEEDLE 29G 12MM157 | XGEVA | |
| VERIFINE PEN NEEDLE 31G 5MM157 | XIFAXAN ORAL TABLET 200 MG, 55 | 0 |
| VERIFINE PEN NEEDLE 31G X 6MM157 | MG | |
| VERIFINE PEN NEEDLE 31G X 8MM157 | XOLAIR | 234 |
| VERIFINE PEN NEEDLE 32G 6MM 157 | XOSPATA | 137 |
| VERIFINE PEN NEEDLE 32G X 4MM157 | | |
| VERIFINE PEN NEEDLE 32G X 5MM157 | | |
| VERIFINE PLUS PEN NDL 31G 5MM 157 | | |
| VERIFINE PLUS PEN NDL 31G 8MM 157 | | |
| VERIFINE PLUS PEN NDL 32G 4MM 157 | | |

| XPOVIO ORAL TABLET 100 | |
|-----------------------------------|------|
| MG/WEEK (50 MG X 2), 40 MG/WEEK | |
| (10 MG X 4), 40 MG/WEEK (20 MG X | |
| 2), 40 MG/WEEK (40 MG X 1), 40MG | |
| TWICE WEEK (40 MG X 2), 60 | |
| MG/WEEK (60 MG X 1), 60MG TWICE | |
| WEEK (120 MG/WEEK), 80 MG/WEEK | |
| (40 MG X 2), 80MG TWICE WEEK (160 | |
| MG/WEEK) | |
| XTANDI ORAL CAPSULE | 110 |
| XTANDI ORAL TABLET 40 MG, 80 MG | Ĵ |
| | 110 |
| XYOSTED | .327 |
| YERVOY | .171 |
| YESINTEK368, | |
| YONSA | 8 |
| YUFLYMA(CF) | 14 |
| YUFLYMA(CF) AI CROHN'S-UC-HS | 14 |
| YUFLYMA(CF) AUTOINJECTOR | 14 |
| ZEJULA ORAL CAPSULE | |
| ZEJULA ORAL TABLET | 219 |
| ZELBORAF | 375 |
| ZIIHERA | 383 |
| ZIRABEV | 54 |
| ZOLADEX | 143 |
| ZTALMY | .135 |
| ZTLIDO | .192 |
| ZURZUVAE ORAL CAPSULE 20 MG, | |
| 25 MG, 30 MG | 387 |
| ZYDELIG | .149 |
| ZYKADIA | 68 |
| ZYNLONTA | 194 |
| ZYNYZ | .268 |